



PAPER • OPEN ACCESS

## Decay of correlations and absence of superfluidity in the disordered Tonks–Girardeau gas

To cite this article: Robert Seiringer and Simone Warzel 2016 *New J. Phys.* **18** 035002

View the [article online](#) for updates and enhancements.

### Related content

- [Localization via fractional moments for models on  \$\mathbb{Z}\$  with single-site potentials of finite support](#)  
A Elgart, M Tautenhahn and I Veseli
- [Ground state asymptotics of a dilute, rotating gas](#)  
Robert Seiringer
- [Dynamical localization for a multi-particle alloy-type model](#)  
Victor Chulaevsky, Anne Boutet de Monvel and Yuri Suhov

### Recent citations

- [Low-Energy Fock-Space Localization for Attractive Hard-Core Particles in Disorder](#)  
Vincent Beaud and Simone Warzel
- [On Polynomial Lieb–Robinson Bounds for the XY Chain in a Decaying Random Field](#)  
Martin Gebert and Marius Lemm
- [Many-body localization for disordered Bosons](#)  
Günter Stolz



## PAPER

## Decay of correlations and absence of superfluidity in the disordered Tonks–Girardeau gas

## OPEN ACCESS

RECEIVED  
21 December 2015REVISED  
26 January 2016ACCEPTED FOR PUBLICATION  
27 January 2016PUBLISHED  
29 February 2016Robert Seiringer<sup>1</sup> and Simone Warzel<sup>2</sup><sup>1</sup> IST Austria, Am Campus 1, A-3400 Klosterneuburg, Austria<sup>2</sup> Zentrum Mathematik, TU München, Boltzmannstr. 3, D-85747 Garching, GermanyE-mail: [robert.seiringer@ist.ac.at](mailto:robert.seiringer@ist.ac.at) and [warzel@ma.tum.de](mailto:warzel@ma.tum.de)**Keywords:** Tonks–Girardeau gas, superfluidity, many-body localization, decay of correlationsOriginal content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/4.0/).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

**Abstract**

We consider the Tonks–Girardeau gas subject to a random external potential. If the disorder is such that the underlying one-particle Hamiltonian displays localization (which is known to be generically the case), we show that there is exponential decay of correlations in the many-body eigenstates. Moreover, there is no Bose–Einstein condensation and no superfluidity, even at zero temperature.

**1. Introduction**

Understanding the various aspects and even the qualitative structure of phase diagrams of interacting many-body systems in the presence of static disorder still poses a big challenge. Basic questions, such as the existence and characterizations of a phase of many-body localized states, remain under debate—even for one-dimensional systems (see [6, 40] and references therein). For bosons, one manifestation of localization is the existence of a glass phase in which the static correlations decay and superfluidity is absent [16]. While such a phase is predicted to exist for strong interactions or strong disorder, for intermediate interaction strength superfluidity is expected to persist at small values of the disorder even in one-dimension [19, 46]. The interest in these questions was renewed due to experimental accessibility of such systems [38] (see also [9, 49] and references therein).

In this context, and in view of the woefully short list of rigorous results on disordered systems with interaction [24, 41], limiting or integrable model systems present a testing ground for numerical works, conjectures and ideas (see [8, 29, 30, 50, 52]). In the bosonic case, the limiting case of hard-core repulsive interaction is such an example: in the lattice set-up this amounts to studying the *XY*-spin Hamiltonian with a random magnetic field, and in the continuum this is the Tonks–Girardeau model with a random potential, which is the topic of the present paper. Such hard-core interactions may actually be realized experimentally [27, 42]—albeit without disorder. Both models can be related to non-interacting fermions in an external random potential. They are not exactly solvable, but nevertheless amenable to rigorous analysis; the difficulty in both cases lies in the non-local dependence of the physical (bosonic) correlation functions on the underlying fermionic correlation functions. In [11] this link was exploited numerically to show that the disorder destroys bosonic quasi-long-range order. For the *XY*-model such a result can be confirmed by rigorous bounds on the correlations of any eigenstate [53] (see also [1, 2, 23, 28] for related and earlier results in this context). The purpose of this paper is to show that such results also apply to the corresponding continuum model. In addition to a proof of the exponential decay of correlations for all eigenstates, we show that the superfluid density (or stiffness) vanishes (exponentially) at zero temperature. Our basic assumption in all these results is the (exponential) localization of the underlying one-particle operator—a property which generically holds true up to arbitrarily large energies in one-dimensional disordered systems [37, 43].

We consider a system of  $N$  bosons with point interactions on a ring with length  $L$ , which we take to be an integer for simplicity. It is described by a many-particle Hamiltonian of the form

$$\mathcal{H}_{L,\omega} = \sum_{j=1}^N (H_{L,\omega}^+)_j + g \sum_{1 \leq j < k \leq N} \delta(x_j - x_k). \quad (1.1)$$

We will be interested in the cases when the one-particle Hamiltonian is given by

$$H_{L,\omega}^{\pm} := -\frac{d^2}{dx^2} + V_{\omega}(x) \tag{1.2}$$

on  $L^2([0, L])$  with periodic boundary conditions (b.c.s). The dependence on  $\omega$ , which will often be omitted from the notation for convenience, indicates the randomness entering the potential landscape. We will assume throughout that the following probabilistic average is finite

$$\sup_n \mathbb{E} \left[ \int_{I_n} |V(x)| dx \right] < \infty, \quad I_n := (n - 1, n], \tag{1.3}$$

where  $\mathbb{E}$  stand for the expectation with respect to  $\omega$ . This ensures, in particular, that  $V_{\omega} \in L^1([0, L])$  and that the one-particle Hamiltonian (1.2) (defined via its quadratic form on the Sobolev space  $H^1[0, L]$ ) is self-adjoint in  $L^2([0, L])$  (with any self-adjoint b.c.s and, in particular, with periodic ones) for almost all  $\omega$ .

In the Tonks–Girardeau limit  $g \rightarrow \infty$ , the bosonic wavefunctions are required to vanish upon particle contact, i.e.,  $\Psi(x_1, \dots, x_N) = 0$  in case  $x_j = x_k$  for some  $j \neq k$ . Any eigenfunction of  $\mathcal{H}_{L,\omega}$  hence takes the form of an eigenfunction of a system of  $N$  non-interacting fermions multiplied by a suitable phase to render it symmetric upon particle exchange [20, 56]. More precisely, let  $H_L^{\pm}$  stand for (1.2) with periodic (+) or anti-periodic (−) b.c.s. If  $\{\varphi_{j,L}^{\pm}\}$  denotes an eigenbasis of  $H_L^{\pm}$  and  $\{j_{\alpha}\}_{\alpha=1}^N$  indexes a subset of  $N$  orthonormal eigenfunctions, then

$$\Psi(x_1, \dots, x_N) = \frac{1}{\sqrt{N!}} \det(\varphi_{j_{\alpha},L}^{\pm}(x_{\beta}))_{\alpha,\beta=1}^N \prod_{1 \leq j < k \leq N} \text{sign}(x_j - x_k) \tag{1.4}$$

is a normalized eigenfunction of the Tonks–Girardeau Hamiltonian provided we choose  $\sharp_N := (-1)^{N+1}$ , i.e., *anti-periodic* b.c. in case  $N$  is even and *periodic* b.c. in case  $N$  is odd (see [34]). In particular, the bosonic ground state of  $\mathcal{H}_L$  corresponds to choosing  $\{j_{\alpha}\}_{\alpha=1}^N$  the  $N$  lowest eigenvalues of  $H_L^{\sharp_N}$ , and its ground state energy  $E_L(N)$  is simply the sum of the lowest  $N$  eigenvalues of  $H_L^{\sharp_N}$ .

We will mainly investigate two quantities of interest:

- (1) The reduced one-particle density matrix  $\gamma_{\Psi}$  corresponding to any eigenstate  $\Psi$  given by (1.4). It is defined through its kernel

$$\gamma_{\Psi}(x, y) := N \int \Psi(x, x_2, \dots, x_N) \overline{\Psi(y, x_2, \dots, x_N)} dx_2 \dots dx_N, \tag{1.5}$$

and satisfies  $0 \leq \gamma_{\Psi} \leq N$  and  $\text{Tr } \gamma_{\Psi} = N$ . Bose–Einstein condensation (BEC) refers to a macroscopic value of the largest eigenvalue  $\|\gamma_{\Psi}\|$  of this operator in the thermodynamic limit ( $N, L \rightarrow \infty$  with  $\frac{N}{L} = \text{const.}$ ) [44] (see also [36]).

It is not hard to show that the reduced one-particle density matrix takes the form of a determinant of an  $(N + 1) \times (N + 1)$  block matrix

$$\gamma_{\Psi}(x, y) = \det \begin{pmatrix} 0 & \varphi_{j_1,L}^{\sharp_N}(x) \cdots \varphi_{j_N,L}^{\sharp_N}(x) \\ \overline{\varphi_{j_1,L}^{\sharp_N}(y)} & \\ \vdots & K_N(x, y) \\ \overline{\varphi_{j_N,L}^{\sharp_N}(y)} & \end{pmatrix}, \tag{1.6}$$

where the  $N \times N$ -submatrix  $K_N(x, y)$  is for all  $x \leq y$  given by the entries

$$\begin{aligned} [K_N(x, y)]_{\alpha,\beta} &:= \delta_{\alpha,\beta} - 2 \int_{[x,y]} \varphi_{j_{\alpha},L}^{\sharp_N}(z) \overline{\varphi_{j_{\beta},L}^{\sharp_N}(z)} dz \\ &= -\delta_{\alpha,\beta} + 2 \int_{[x,y]^c} \varphi_{j_{\alpha},L}^{\sharp_N}(z) \overline{\varphi_{j_{\beta},L}^{\sharp_N}(z)} dz. \end{aligned} \tag{1.7}$$

Introducing the projection  $P_N = \sum_{\alpha} |\varphi_{j_{\alpha},L}^{\sharp_N}\rangle \langle \varphi_{j_{\alpha},L}^{\sharp_N}|$  onto the eigenfunctions entering the state  $\Psi$ , we may write  $K_N(x, y) = P_N - 2P_N 1_{[x,y]} P_N$  as an operator identity on  $P_N L^2([0, L])$ . In this manner, one easily sees that  $\gamma_{\Psi}$  only depends on the projection  $P_N$ , as a change of basis corresponds to a unitary transformation of the matrix in (1.6) which leaves the determinant invariant.

- (2) The superfluid density (or: stiffness) measures the extent to which the ground state energy of the Tonks–Girardeau Hamiltonian increases as one twists the b.c.s [15], i.e., when the wave-functions are required to pick up a phase  $e^{i\theta}$  as one particle moves around the ring,  $\Psi(x_1, \dots, x_j + L, \dots, x_N) = e^{i\theta} \Psi(x_1, \dots, x_j, \dots, x_N)$ . In other words, the superfluid density  $\rho_s$  is defined via the ground-state energy shift  $E_L(N, \theta) \approx E_L(N, 0) + \theta^2 \rho_s / L$  for small  $\theta$ . To give a precise definition in the thermodynamic limit, we find

it more convenient to work in a grand-canonical picture where the particle number is determined by a fixed chemical potential  $\mu$ ; i.e.,  $\mu$  is chosen independently of  $N$  and  $\omega$ .

For given  $\mu \in \mathbb{R}$ , let  $N_\mu^\pm := \text{Tr } 1_{(-\infty, \mu]}(H_L^\pm)$  denote the number of eigenstates of  $H_L^\pm$  below  $\mu$  and set  $N_\mu := \min\{N_\mu^+, N_\mu^-\}$ . With  $E_L(N, \theta)$  the ground state energy of the Tonks–Girardeau Hamiltonian with twisted b.c.s, the superfluid density is defined as

$$\rho_s := \limsup_{\theta \rightarrow 0} \frac{1}{\theta^2} \limsup_{L \rightarrow \infty} L(E_L(N_\mu, \theta) - E_L(N_\mu, 0)). \tag{1.8}$$

We note that it follows from the diamagnetic inequality [35, theorem 7.21] that  $E_L(N, \theta) \geq E_L(N, 0)$  for any  $\theta$  and  $N$ , hence  $\rho_s \geq 0$ .

With our definition of  $N_\mu$ , the ground state energy  $E_L(N_\mu, 0)$  has the following convenient representation in terms of  $H_L^\pm$ . With  $\sharp_\mu := \sharp_{N_\mu} = (-1)^{N_\mu+1}$ , we have

$$E_L(N_\mu, 0) = \mu N_\mu - \text{Tr}[H_L^{\sharp_\mu} - \mu]_- = \sum_{j: E_{j,L}^{\sharp_\mu} \leq \mu} E_{j,L}^{\sharp_\mu}, \tag{1.9}$$

where the  $\{E_{j,L}^\pm\}$  denote the eigenvalues of  $H_L^\pm$ , and  $[\cdot]_- := -\min\{0, \cdot\}$  denotes the negative part. In other words, for any  $\mu \in \mathbb{R}$ ,  $H_L^{\sharp_\mu}$  has exactly  $N_\mu$  eigenvalues below  $\mu$ . This is a consequence of the fact that  $N_\mu = N_\mu^{\sharp_\mu}$ , which, in turn, follows from the interlacing property  $E_{j,L}^+ < E_{j,L}^-$  for  $j$  odd, and  $E_{j,L}^+ > E_{j,L}^-$  for  $j$  even (see [12, theorem 2.3.1] or [47, theorem XIII.89]).

## 2. Results

### 2.1. Localization hypothesis and first consequences

We will assume that the one-particle operator  $H_{L,\omega}^\pm$  in (1.2), for both periodic (+) and anti-periodic (−) b.c., exhibits (sub-)exponential Anderson localization with some exponential parameter  $\xi \in (0, 1]$  and localization length  $\ell < \infty$  in the energy regimes of interest. To be more specific, we consider the eigenfunction correlator

$$Q_L^\pm(n, m; J; \omega) := \sum_{j, E_{j,L}^\pm \in J} \Phi_{j,L}^\pm(n; \omega) \Phi_{j,L}^\pm(m; \omega) \tag{2.1}$$

corresponding to some energy regime  $J \subset \mathbb{R}$ . Here

$$\Phi_{j,L}^\pm(n; \omega) := \left( \int_{I_n} |\varphi_{j,L}^\pm(x; \omega)|^2 dx \right)^{\frac{1}{2}} \tag{2.2}$$

quantifies the probability for the  $j$ th eigenfunction to be present on a basic interval of unit length. We shall tacitly assume that the complete orthonormal set of eigenfunctions of  $H_{L,\omega}^\pm$  is jointly measurable in  $(x, \omega)$ . (In case of degeneracy, which generically is believed to be absent with probability one, this in particular requires the choice of a proper labelling of eigenfunctions.)

*Localization hypothesis eigenfunction correlator localization (ECL) on  $J$ :* There exist  $C, \ell \in (0, \infty)$  and  $\xi \in (0, 1]$  such that for all  $1 \leq n, m \leq L$  and all  $L \in \mathbb{N}$

$$\mathbb{E}[Q_L^\pm(n, m; J)] \leq C \exp\left(-\frac{\text{dist}(n, m)^\xi}{\ell^\xi}\right), \tag{2.3}$$

where  $\text{dist}(\cdot, \cdot)$  denotes the Euclidean distance on the (one-dimensional) torus.

In the theory of (one-particle) random operators, the condition (ECL) is both strong and convenient: it ensures localization in both the spectral sense (i.e. only pure-point spectrum in  $J$  with (sub-)exponentially decaying eigenfunctions) as well as in the strong dynamical sense that

$$\mathbb{E}[\sup_t \|1_{I_n} e^{-itH_L^\pm} P_J(H_L^\pm) 1_{I_m}\|_1] \leq C \exp\left(-\frac{\text{dist}(n, m)^\xi}{\ell^\xi}\right). \tag{2.4}$$

Here  $P_J(H_L^\pm)$  denotes the spectral projection of  $H_L^\pm$  onto the energy regime  $J$  and  $\|\cdot\|_1 := \text{Tr}|\cdot|$  is the trace norm. ECL is established for a large class of single-particle random Schrödinger operators by means of either the continuum fractional-moment method [4] (which is based on [3]) or via the bootstrap multi-scale analysis [17, 18] (which is based on [14]). In our one-dimensional set-up, localization is expected to hold generically at all energies. In particular, (ECL) will hold for  $J = (-\infty, \mu]$  with  $\xi = 1$  and some localization length  $\ell = \ell_\mu$  which depends on  $\mu \in \mathbb{R}$  only. This has been established in the following specific models:

- Early on in the history of localization proofs [21] for random potentials of the form  $V_\omega(x) = F(b_x(\omega))$  with  $b_x(\omega)$  a Brownian motion on a compact Riemannian manifold  $M$  and  $F: M \rightarrow \mathbb{R}$  a smooth Morse function with  $\min_M F = 0$ .
- For homogeneous *Gaussian random potentials*, i.e.,  $\mathbb{E}[V(x)] = 0$  with covariance function  $C(x - y) := \mathbb{E}[V(x)V(y)]$ , which admits the representation  $C(x) = \int \gamma(x + y)\gamma(y)dy$  in terms of a non-negative, compactly supported function  $\gamma$ , which is uniformly Hölder continuous, i.e., there is  $s \in (0, 1]$  and  $a < \infty$  such that  $|\gamma(x + y) - \gamma(x)| \leq a|y|^s$  for all  $x$  and all  $y > 0$  sufficiently small (see [13, 55]).
- For *alloy-type random potentials*  $V_\omega(x) = W(x) + \sum_{j \in \mathbb{Z}} \omega_j U(x - j)$  with independent and identically distributed random variables  $(\omega_j)_{j \in \mathbb{Z}}$  whose distribution is absolutely continuous with a bounded density, i.e.  $\mathbb{P}(\omega_j \in dv) = \varrho(v)dv$  with some  $\varrho \in L^\infty(\mathbb{R}) \cap L^1(\mathbb{R})$  of compact support. The term  $W$  serves as a non-random, bounded, 1-periodic background potential and the single-site potential  $U$  is assumed to satisfy  $c_I(x) \leq U(x) \leq C1_{[0,1]}(x)$  for some  $c, C \in (0, \infty)$  and a non-trivial sub-interval  $I \subset [0, 1]$  (see [22]).

ECL implies the (sub-)exponential localization of eigenfunctions about some random localization center. More precisely, it implies what is called semi-uniform localization of eigenfunctions (SULE). For the definition of the latter it is convenient to fix a weight function  $g_L: \{1, \dots, L\} \rightarrow [1, \infty)$  with the property  $\sum_{\alpha=1}^L g_L(\alpha)^{-1} = 1$ . A specific choice, which we will adopt below, is  $g_L(\alpha) = L$ .

*Localization hypothesis (SULE) on J:* There exist  $\ell \in (0, \infty)$ ,  $\xi \in (0, 1]$  and, for every  $L \in \mathbb{N}$ , an amplitude  $A_{L,\omega} \geq 0$  that is uniformly integrable, i.e.

$$\sup_{L \in \mathbb{N}} \mathbb{E}[A_L] < \infty, \tag{2.5}$$

such that for every eigenfunction  $\varphi_{j,L}^\pm$  of  $H_{L,\omega}^\pm$  with eigenvalue  $E_{j,L}^\pm \in J$  there is some  $\gamma_{j,L,\omega}^\pm \in \mathbb{N}$  such that for all  $n$ :

$$\Phi_{j,L}^\pm(n; \omega) \leq A_{L,\omega} g_L(\gamma_{j,L,\omega}^\pm)^{3/2} \exp\left(-\frac{\text{dist}(n, \gamma_{j,L,\omega}^\pm)^\xi}{\ell^\xi}\right). \tag{2.6}$$

The points  $\gamma_{j,L,\omega}^\pm$  play the role of localization centers. However, they need not coincide with the location of the maxima of  $\Phi_{j,L}^\pm(n; \omega)$ . The length  $\ell$  is non-random and coincides with the minimum of all localization length at energies in  $J$ . At first sight, the role of the function  $g_L$  might be puzzling and one may be tempted to drop the factor  $g_L^{3/2}$  on the right side of (2.6). This, however, is known to be wrong [10]. If assumption (ECL) holds for an energy regime  $J$ , then (SULE) holds with any weight function  $g_L$  for the same energy regime (but possibly with a slightly reduced localization length), see [5, chapter 7].

As explained in the introduction, every fermionic many-body state  $\phi(x_1, \dots, x_N)$ , which is either periodic or antiperiodic depending on whether  $N$  is odd or even, gives rise to the periodic bosonic many-body state  $\Psi(x_1, \dots, x_N) = \phi(x_1, \dots, x_N) \prod_{1 \leq j < k \leq N} \text{sign}(x_j - x_k)$ . In the Tonks–Girardeau limit, the dynamics of such a state is given in terms of the dynamics of free fermions, i.e.,  $\phi_t(x_1, \dots, x_N) = \exp\left(-it \sum_{j=1}^N (H_L^{\#N})_j\right) \phi(x_1, \dots, x_N)$ . While the bosonic one-particle density matrix  $\gamma_{\Psi_t}$  of this state does not coincide with the fermionic one, given by  $\Gamma_{\phi_t}$  (which is defined as in (1.5) with  $\Psi$  replaced by  $\phi_t$ ), their diagonals agree, i.e., the bosonic and fermionic densities are equal:

$$\varrho_t(x) := \gamma_{\Psi_t}(x, x) = \Gamma_{\phi_t}(x, x) = (e^{-itH_L^{\#N}} \Gamma_\phi e^{itH_L^{\#N}})(x, x). \tag{2.7}$$

Dynamical localization for free fermions, in the form (2.4), then immediately entails the following result for any many-body eigenstate. The bound (2.8) is a manifestation of many-body localization for the model of interacting bosons considered here.

**Proposition 2.1.** *If the range of  $\Gamma_\phi$  at  $t = 0$  falls within a regime of dynamical localization, i.e.*

$\Gamma_\phi = P_J(H_L^{\#N})\Gamma_\phi = \Gamma_\phi P_J(H_L^{\#N})$  for some  $J \subset \mathbb{R}$  for which (2.4) holds, then there exists an  $A \in (0, \infty)$  which is independent of  $L$  and  $N$  such that:

(1) *the total number of particles on any subset  $I \subset [0, L]$  changes on average by order one only:*

$$\mathbb{E} \left[ \sup_{t \in \mathbb{R}} \left| \int_I \varrho_t(x) dx - \int_I \varrho_0(x) dx \right| \right] \leq A \quad \text{for all } I \subset [0, L], \tag{2.8}$$

(2) for any pair of subsets  $I \subset K \subset [0, L]$ :

$$\mathbb{E} \left[ \sup_{t \in \mathbb{R}} \int_I \varrho_t(x) dx \right] \leq \mathbb{E} \left[ \int_K \varrho_0(x) dx \right] + A \exp \left( - \frac{\text{dist}(I, K^c)^\xi}{\ell^\xi} \right). \quad (2.9)$$

The (simple) proof of this proposition will be given in the appendix. Both statements are expressions of non-ergodic behavior of the localized system: one may prepare the system initially in a state which exhibits a step-like profile in its density, i.e., some positive averaged density in one half ( $I$ ) and another one in the other half ( $I^c$ ). In such a situation, (2.8) states that the step-like profile remains for arbitrarily long times with only a finite number of particles crossing on average.

The second bound is relevant for experiments in which the bosons are initially trapped around some location (such that  $\int_K \varrho_0(x) dx \approx 0$ ) and then released from the trap at  $t = 0$ . The localization bound (2.9) then guarantees that the total number of particles will remain small on average away from the initial location, uniformly in time (confirming numerical simulations in [45].) A related bound for the XY-model can be found in [2, theorem 1.1].

## 2.2. Decay of correlations and absence of BEC

Our first non-trivial consequence of one-particle localization concerns a strong version of absence of off-diagonal long-range order (ODLRO).

**Theorem 2.2.** *Let  $\Psi$  be any many-particle eigenstate of the form (1.4) which is composed of a selection of one-particle states  $\{\varphi_{j_\alpha, L}^{\#N}\}_{\alpha=1}^N$  corresponding to an energy regime  $J$ . If condition (ECL) holds for  $J$ , then there exist  $A \in (0, \infty)$  independent of  $L$  and  $N$  such that*

$$\mathbb{E}[\|1_n \gamma_\Psi 1_m\|_2^\sigma] \leq A \exp \left( - \frac{2}{3} (1 - \sigma) \frac{\text{dist}(n, m)^\xi}{(2\ell)^\xi} \right) \quad (2.10)$$

for all  $1 \leq n, m \leq L$  and all  $2/5 \leq \sigma < 1$ . Here  $\|\cdot\|_2$  denotes the Hilbert–Schmidt-norm on  $L^2([0, L])$ .

A proof of this theorem, as well as of the subsequent corollary, will be given in section 3. The proof shows that the result can easily be extended in various directions, and is not restricted to eigenstates of the many-particle Hamiltonian. It applies, e.g., to general states of the form (1.4) as long as the one-particle functions  $\varphi_j$  are suitably localized, and is thus also relevant in time-dependent situations as in [48].

Absence of BEC is not immediately implied by the absence of ODLRO, since our assumptions on the system allow for unbounded fluctuations of the density. We therefore need a mild additional assumption on these fluctuations in order to reach such a conclusion.

**Corollary 2.3.** *Given the assumptions of theorem 2.2, assume additionally that for  $p > 2$*

$$\sup_{n, L} \mathbb{E}[(\text{Tr } 1_n P_J (H_L^\pm))^p] < \infty. \quad (2.11)$$

*Then for any sequence  $\Psi$  of eigenstates composed of one-particle states  $(\varphi_{j_\alpha, L}^{\#N})_{\alpha=1, \dots, N}$  whose energies fall into a regime  $J$  of (ECL), the almost-sure convergence*

$$\lim_{L \rightarrow \infty} \frac{\|\gamma_\Psi\|}{L} = 0 \quad (2.12)$$

*holds for any  $\frac{2}{p} < r \leq 1$ .*

Note that the convergence (2.12) is independent of the choice of  $N$ , which is allowed to depend on  $L$  and  $\omega$ . Typically one is interested in the case that  $N \approx \text{const. } L$  as  $L \rightarrow \infty$ . The subsequent proof (specifically equation (3.30)) also shows that in case (2.11) holds with  $p > 1$  the (averaged) momentum distribution associated with the state  $\Psi$ ,

$$n(k) := \frac{1}{L} \iint e^{ik(x-y)} \mathbb{E}[\gamma_\Psi(x, y)] dx dy, \quad (2.13)$$

remains uniformly bounded, since  $|n(k)| \leq \sup_L \frac{1}{L} \sum_{n, m} \mathbb{E}[\|1_n \gamma_\Psi 1_m\|_2] < \infty$ . (In particular,  $n(0) < \infty$ . At large values  $|k| \rightarrow \infty$ , one expects an algebraic fall-off  $n(k) \sim k^{-4}$  due to the hard-core repulsion [7]). This is consistent with numerical predictions in [11].

A simple sufficient condition for (2.11) to hold for any value  $p > 2$  (and hence for the validity of (2.12) for any  $r > 0$ ) and  $\mathbb{Z}$ -homogeneous random potentials in case  $J \subset (-\infty, \mu]$  is the existence of an exponential moment

$$\int_{I_t} \mathbb{E}[e^{-tV(x)}] dx < \infty \tag{2.14}$$

for some  $t > 0$ . (A derivation of this statement starts from the estimate  $\text{Tr } 1_{I_n} P_{(-\infty, \mu]}(H_L^\pm) \leq e^{t\mu} \text{Tr } 1_{I_n} e^{-tH_L^\pm}$  and proceeds through standard semigroup bounds, see [43, chapter II 5].) This is clearly satisfied for the models listed above. (Alternatively, one may proceed through resolvent techniques as in [43, chapter II 5] to show that the finiteness of  $\sup_n \int_{I_n} \mathbb{E}[|V(x)|^{2p}] dx$  is sufficient for (2.11) to hold.)

The absence of ODLRO and BEC is not a consequence of the disorder alone: In case  $V = 0$ , Lenard [31] showed that the reduced density matrix of the ground state wave function  $\Psi$  behaves as  $\gamma_\Psi(x, y) \sim |x - y|^{-1/2}$  for large  $|x - y|$ . This slow fall off is sometimes referred as quasi-long range order and causes of the order of  $\sqrt{N}$  particles to quasi-condense into the zero mode.

The above corollary (with  $r < 1/2$ ) shows that localization decreases the rate of quasi-condensed particles in comparison to the free case ( $V = 0$ ). This should not be taken for granted as a comparison with the non-interacting case shows. For *non-interacting* bosons in a non-negative Poisson random potential, Luttinger together with Kac [25, 26] and Sy [39] noted that that critical dimension  $d$  for the occurrence of BEC is lowered to  $d = 1$ . (A rigorous version of their analysis is contained in [32, 33] and the basic mechanism also applies to alloy-type random potentials.) The occurrence of BEC in an ideal Bose gas even at positive temperature is due to the behavior of the density of states near the bottom of the one-particle energy spectrum. The latter is severely suppressed due to the occurrence of Lifshitz tails, which causes a macroscopic fraction of the particles to condense into modes whose energy vanishes in the thermodynamic limit. Since Anderson localization is known for such models, our results imply that the interactions destroy BEC, and the corresponding Tonks–Girardeau model shows no BEC even at zero temperature.

### 2.3. Absence of superfluidity

In the absence of an external random potential ( $V = 0$ ), it is well-known that the superfluid density (1.8) coincides with the total density at chemical potential  $\mu$ :

$$\rho_s = \lim_{L \rightarrow \infty} \frac{1}{L} N_\mu = \frac{\sqrt{[\mu]_+}}{2\pi^2}. \tag{2.15}$$

In particular, it is strictly positive for all  $\mu > 0$ . This changes drastically in the regime of localization.

**Theorem 2.4.** *If condition (SULE) holds (with  $g_L(\alpha) = L$ ) for the energy regime  $(-\infty, \mu]$ , then for any  $\theta > 0$  and almost surely:*

$$\limsup_{L \rightarrow \infty} L(E_L(N_\mu, \theta) - E_L(N_\mu, 0)) = 0. \tag{2.16}$$

*As a consequence, the superfluid density (1.8) is zero almost surely.*

Our result implies that generically a disordered Bose gas in one-dimension in the Tonks–Girardeau regime shows no superfluidity and no BEC even at zero temperature. This statement concerns the usual thermodynamic limit. We note that other limiting regimes are possible, corresponding to mean-field type interactions, where both BEC and superfluidity can prevail at zero temperature [30, 50] (see also [8, 29, 54] for related results). The proof of theorem 2.4 will be given in section 4.

## 3. Proof of decay of correlations

This section is devoted to the proofs of theorem 2.2 and corollary 2.3. Since  $N$  is kept fixed in theorem 2.2, we will drop the superscript  $\sharp_N$  on the eigenfunctions  $\{\varphi_{j,L}^{\sharp_N}\}$  of  $H_L^{\sharp_N}$  (as well as their dependence on  $\omega$ ) for ease of notation.

Using the Laplace formula, the determinantal expression (1.6) for the kernel of the one-particle reduced density matrix can be recast as

$$\gamma_\Psi(x, y) = \sum_{\alpha, \beta} \varphi_{j_\alpha, L}(x) \overline{\varphi_{j_\beta, L}(y)} [\text{adj } K_N(x, y)]_{\beta, \alpha} =: \langle \varphi(y), \text{adj } K_N(x, y) \varphi(x) \rangle, \tag{3.1}$$

where the last inner product is in  $\mathbb{C}^N$  and the adjugate matrix of  $K_N(x, y)$  is the matrix of cofactors (up to signs), i.e.



$$[\text{adj } K_N(x, y)]_{\beta, \alpha} = \det \begin{pmatrix} 0 & e_\alpha^T \\ e_\beta & K_N(x, y) \end{pmatrix} = (-1)^{\alpha+\beta} \det[K_N(x, y)]_{\hat{\alpha}, \hat{\beta}}. \tag{3.2}$$

Here  $\{e_\alpha\}$  denote the unit vectors and the hats indicate the deletion of row  $\beta$  and column  $\alpha$  from  $K_N(x, y)$ . Some key properties are summarized in the following:

- (1) In case  $x = y$ ,  $K_N(x, x) = \text{adj } K_N(x, x)$  equals the identity matrix.
- (2) Since  $K_N(x, y) = P_N - 2P_N \mathbb{1}_{[x,y]} P_N$ , the inequality  $-P_N \leq K_N(x, y) \leq P_N$  holds, and hence we arrive at the bound  $\|K_N(x, y)\| \leq 1$  on the operator norm.
- (3) Since the adjugate matrix of any hermitian  $N \times N$  matrix is hermitian with eigenvalues given by the products of  $N - 1$  disjoint eigenvalues of the matrix, the norm bound

$$\|\text{adj } K_N(x, y)\| \leq 1 \tag{3.3}$$

is an immediate consequence of  $\|K_N(x, y)\| \leq 1$ .

Our basic strategy for an estimate of (3.1) is to split the summation depending on whether the eigenstates live predominantly to the right or left of the midpoint of  $n$  and  $m$ . To do so, we will suppose without loss of generality  $1 \leq n < m \leq L/2$  and abbreviate by

$$M := \lfloor (m + n)/2 \rfloor \tag{3.4}$$

the midpoint between  $n$  and  $m$  (or between  $n$  and  $m - 1$  if  $(m + n)/2$  is not an integer). This midpoint introduces a left/right partition of the system according to which we may sort the eigenstates:

$$\mathcal{L} := \left\{ \alpha \mid \int_0^M |\varphi_{j_\alpha, L}(\xi)|^2 > \frac{1}{2} \right\}, \quad \mathcal{R} := \left\{ \alpha \mid \int_M^L |\varphi_{j_\alpha, L}(\xi)|^2 \geq \frac{1}{2} \right\}. \tag{3.5}$$

The normalization of eigenstates implies that  $\mathcal{L}$  and  $\mathcal{R}$  constitute a disjoint partition of the finite index set  $\{1, \dots, N\}$ . Writing the vectors on the right side of (3.1) accordingly as  $\varphi(x) = \varphi_{\mathcal{L}}(x) + \varphi_{\mathcal{R}}(x)$  (and similarly for  $\varphi(y)$ ) we may split the sum (3.1) into three parts,  $\gamma_\Psi(x, y) = \gamma_\Psi^{(1)}(x, y) + \gamma_\Psi^{(2)}(x, y) + \gamma_\Psi^{(3)}(x, y)$ , with

$$\begin{aligned} \gamma_\Psi^{(1)}(x, y) &:= \langle \varphi_{\mathcal{L}}(y), \text{adj } K_N(x, y) \varphi(x) \rangle, \\ \gamma_\Psi^{(2)}(x, y) &:= \langle \varphi_{\mathcal{R}}(y), \text{adj } K_N(x, y) \varphi_{\mathcal{R}}(x) \rangle \\ \gamma_\Psi^{(3)}(x, y) &:= \langle \varphi_{\mathcal{R}}(y), \text{adj } K_N(x, y) \varphi_{\mathcal{L}}(x) \rangle. \end{aligned} \tag{3.6}$$

The Hilbert–Schmidt norms of these contributions are estimated separately. We start with the first two terms.

**Lemma 3.1.** For all  $m > n$ :

$$\begin{aligned} \| \mathbb{1}_{I_n} \gamma_\Psi^{(1)} \mathbb{1}_{I_m} \|_2 &\leq \sqrt{2 N(I_n)} \sum_{k=1}^M Q_L(k, m; J) \\ \| \mathbb{1}_{I_n} \gamma_\Psi^{(2)} \mathbb{1}_{I_m} \|_2 &\leq \sqrt{2 N(I_m)} \sum_{k=M+1}^L Q_L(n, k; J), \end{aligned} \tag{3.7}$$

where  $M$  was defined in (3.4) and  $N(I_k) := \int_{I_k} P_N(x, x) dx$  denotes the local particle number in  $I_k$ .

**Proof.** The Cauchy–Schwarz inequality in  $\mathbb{C}^N$  and (3.3) imply

$$|\gamma_\Psi^{(1)}(x, y)|^2 \leq \|\varphi_{\mathcal{L}}(y)\|^2 \|\varphi(x)\|^2 \|\text{adj } K_N(x, y)\|^2 \leq P_N(x, x) \sum_{\alpha \in \mathcal{L}} |\varphi_{j_\alpha, L}(y)|^2. \tag{3.8}$$

Integration over  $x \in I_n$  and  $y \in I_m$  yields the bound

$$\| \mathbb{1}_{I_n} \gamma_\Psi^{(1)} \mathbb{1}_{I_m} \|_2^2 \leq N(I_n) \sum_{\alpha \in \mathcal{L}} \Phi_{j_\alpha, L}(m)^2. \tag{3.9}$$

The last term may be estimated using the fact that eigenfunctions corresponding to  $\alpha \in \mathcal{L}$  predominantly live on the left:

$$\sum_{\alpha \in \mathcal{L}} \Phi_{j_\alpha, L}(m)^2 \leq 2 \sum_{k=1}^M \sum_{\alpha \in \mathcal{L}} \Phi_{j_\alpha, L}(k)^2 \Phi_{j_\alpha, L}(m)^2 \leq 2 \sum_{k=1}^M Q_L(k, m; J)^2. \tag{3.10}$$



Finally, we use that

$$\sum_{k=1}^M Q_L(k, m; J)^2 \leq \left( \sum_{k=1}^M Q_L(k, m; J) \right)^2. \tag{3.11}$$

This completes the proof of the first inequality. A proof of the second inequality proceeds analogously with the roles of  $\mathcal{L}$  and  $\mathcal{R}$  interchanged.  $\square$

Both terms are thus (sub-)exponentially small in the distance between  $n$  and  $m$  provided the eigenfunction correlator decays accordingly. To establish this for the third term we employ the technique developed in [53] for estimates on certain structured determinants.

**Lemma 3.2.** For all  $m > n$ :

$$\|1_{I_n} \gamma_{\Psi}^{(3)} 1_{I_m}\|_2 \leq 2\sqrt{2e N(I_n) N(I_m)} \sum'_{(k,l)} \sqrt{N(I_k)} Q_L(k, l; J). \tag{3.12}$$

Here we have abbreviated  $\sum'_{(k,l)} := \left[ \sum_{k=1}^n \sum_{l=M+1}^L + \sum_{k=m}^L \sum_{l=1}^M \right]$ .

**Proof.** Using the definition of the adjugate matrix, we rewrite the inner product again as a determinant of a block matrix of the following form:

$$\gamma_{\Psi}^{(3)}(x, y) = \det \begin{pmatrix} 0 & \varphi_{\mathcal{L}}(x)^T & 0 \\ 0 & \tilde{K}_N(x, y) & \\ \frac{\varphi_{\mathcal{R}}(y)}{\varphi_{\mathcal{R}}(y)} & & \end{pmatrix} \tag{3.13}$$

where  $\tilde{K}_N(x, y) = V^T K_N(x, y) V$  for a permutation matrix  $V$  that permutes the indices such that the indices in  $\mathcal{L}$  correspond to the first  $|\mathcal{L}|$  rows and columns, and the ones in the  $\mathcal{R}$  to the last  $|\mathcal{R}| = N - |\mathcal{L}|$  ones. Note that  $\|\tilde{K}_N(x, y)\| \leq 1$ . Hence we can apply the following estimate on the determinant, which is a simple variant of the bound in theorem 3.1 in [53].

**Lemma 3.3.** Let  $v \in \mathbb{C}^p$ ,  $w \in \mathbb{C}^q$ , and let  $K = \begin{pmatrix} A & B \\ C & D \end{pmatrix}$  be a  $(p + q) \times (p + q)$  matrix with  $\|K\| \leq 1$ . Then

$$\left| \det \begin{pmatrix} 0 & v^T & 0 \\ 0 & A & B \\ w & C & D \end{pmatrix} \right| \leq \sqrt{e} \|v\| \|w\| \|B\|. \tag{3.14}$$

**Proof.** By linearity we may assume that  $\|w\| = 1$ . As in [53], we first apply a unitary operator  $U$  on  $\mathbb{C}^p$  that takes  $v$  into the vector  $(0, \dots, 0, \|v\|)$ . Moreover, we can find another unitary  $V$  on  $\mathbb{C}^p$  such that  $V A U^T$  is upper triangular, i.e., all entries below the diagonal are zero. The left side of (3.14) is equal to the absolute value of the determinant of

$$\begin{pmatrix} 1 & 0 & 0 \\ 0 & V & 0 \\ 0 & 0 & 1_q \end{pmatrix} \begin{pmatrix} 0 & v^T & 0 \\ 0 & A & B \\ w & C & D \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & U^T & 0 \\ 0 & 0 & 1_q \end{pmatrix} = \begin{pmatrix} 0 & (Uv)^T & 0 \\ 0 & V A U^T & VB \\ w & C U^T & D \end{pmatrix} =: M. \tag{3.15}$$

To estimate it, we use Hadamard’s bound, which states that the determinant of a matrix is bounded by the product of the norms of the row vectors. Before we apply this bound, we perform one more operation that leaves the determinant invariant, namely we subtract  $s$  times the  $(p + 1)$ th row from the first row, for some  $s \in \mathbb{C}$ . Let  $\alpha$  denote the lower right entry of  $V A U^T$  (i.e., the only non-zero entry in the  $p$ th row of  $V A U^T$ ). Using the fact that the norm of a row vector of a square matrix can never exceed the norm of the matrix, we then obtain

$$|\det M| \leq \sqrt{|\|v\| - s\alpha|^2 + |s|^2 \|B\|^2} \sqrt{|\alpha|^2 + \|B\|^2} \prod_{\alpha=1}^q \sqrt{1 + |w_{\alpha}|^2}. \tag{3.16}$$

The first factor on the right side bounds the norm of the first row, the second one the  $(p + 1)$ th row, and the last factors the rows  $p + 2, \dots, p + q + 1$ . The norms of the other rows  $2, \dots, p$  are bounded by one. Since

$$\prod_{\alpha=1}^q \sqrt{1 + |w_{\alpha}|^2} = \exp \left( \frac{1}{2} \sum_{\alpha=1}^q \ln(1 + |w_{\alpha}|^2) \right) \leq \exp \left( \frac{1}{2} \sum_{\alpha=1}^q |w_{\alpha}|^2 \right) = \sqrt{e}, \tag{3.17}$$

the choice  $s = \bar{\alpha} \|v\| (|\alpha|^2 + \|B\|^2)^{-1}$  leads to the desired bound (3.14).  $\square$

An application of lemma 3.3 to (3.13) leads to the bound

$$|\gamma_{\Psi}^{(3)}(x, y)| \leq \sqrt{e} \|\varphi_{\mathcal{R}}(y)\| \|\varphi_{\mathcal{L}}(x)\| \|B\|, \tag{3.18}$$

where  $B$  is the  $|\mathcal{L}| \times |\mathcal{R}|$  matrix with entries  $\alpha \in \mathcal{L}, \beta \in \mathcal{R}$  given by

$$B_{\alpha,\beta} := [K_N(x, y)]_{\alpha,\beta} = 2 \sum_{k=1}^L \int_{I_k \cap [x,y]^c} \varphi_{j_{\alpha,L}}(z) \overline{\varphi_{j_{\beta,L}}(z)} dz. \tag{3.19}$$

Here the equality results from (1.7). The operator norm of  $B$  is estimated in terms of its Frobenius norm,  $\|B\| \leq \|B\|_2$ , which in turn is bounded from above using Minkowski’s inequality as follows. For  $x \in I_n, y \in I_m$ ,

$$\begin{aligned} \|B\|_2 &= \left( \sum_{\substack{\alpha \in \mathcal{L} \\ \beta \in \mathcal{R}}} |B_{\alpha,\beta}|^2 \right)^{\frac{1}{2}} \leq 2 \sum_{k=1}^L \left( \sum_{\substack{\alpha \in \mathcal{L} \\ \beta \in \mathcal{R}}} \left| \int_{I_k \cap [x,y]^c} \varphi_{j_{\alpha,L}}(z) \overline{\varphi_{j_{\beta,L}}(z)} dz \right|^2 \right)^{\frac{1}{2}} \\ &\leq 2 \left( \sum_{k=1}^n + \sum_{k=m}^L \right) \left( \sum_{\alpha \in \mathcal{L}} \Phi_{j_{\alpha,L}}(k)^2 \sum_{\beta \in \mathcal{R}} \Phi_{j_{\beta,L}}(k)^2 \right)^{\frac{1}{2}} \\ &\leq 2 \sum_{k=1}^n \sqrt{N(I_k)} \left( 2 \sum_{l=M+1}^L Q_L(k, l; J)^2 \right)^{\frac{1}{2}} + 2 \sum_{k=m}^L \sqrt{N(I_k)} \left( 2 \sum_{l=1}^M Q_L(k, l; J)^2 \right)^{\frac{1}{2}} \\ &\leq 2\sqrt{2} \sum'_{(k,l)} \sqrt{N(I_k)} Q_L(k, l; J). \end{aligned} \tag{3.20}$$

The penultimate inequality derives from (3.10) (and its analog for  $\mathcal{R}$ ). After inserting this bound in (3.18) and integrating its square over  $x \in I_n$  and  $y \in I_m$ , we obtain the claimed bound (3.12).  $\square$

We may now conclude the proof of our first main result.

**Proof of theorem 2.2.** We start by noting that  $\mathbb{E}[\|1_{I_n} \gamma_{\Psi} 1_{I_m}\|_2]$  is uniformly bounded. In fact, the Cauchy–Schwarz inequality and the fact that the Hilbert–Schmidt norm is dominated by the trace norm lead to

$$\|1_{I_n} \gamma_{\Psi} 1_{I_m}\|_2 \leq (\|1_{I_n} \gamma_{\Psi} 1_{I_n}\|_2 \|1_{I_m} \gamma_{\Psi} 1_{I_m}\|_2)^{1/2} \leq \sqrt{N(I_n)N(I_m)}. \tag{3.21}$$

Moreover, by the Cauchy–Schwarz inequality for the expectation value

$$\mathbb{E}[\sqrt{N(I_n)N(I_m)}] \leq (\mathbb{E}[N(I_n)] \mathbb{E}[N(I_m)])^{1/2}. \tag{3.22}$$

In turn, the average local particle number is uniformly bounded by assumption:

$$\sup_{L,n} \mathbb{E}[N(I_n)] = \sup_{L,n} \mathbb{E}[Q_L(n, n; J)] \leq C. \tag{3.23}$$

We may therefore assume without loss of generality that  $m > n$ . We may also assume that  $1 \leq n < m \leq L/2$ ; the general case then follows by a simple relabeling. We first proof the assertion for  $\sigma = \frac{2}{5}$ . Since

$$\mathbb{E} \left[ \|1_{I_n} \gamma_{\Psi} 1_{I_m}\|_2^{\frac{2}{5}} \right] \leq \sum_{i=1}^3 \mathbb{E} \left[ \|1_{I_n} \gamma_{\Psi}^{(i)} 1_{I_m}\|_2^{\frac{2}{5}} \right] \tag{3.24}$$

we can treat the three contributions to  $\gamma_{\Psi}$  separately. For the first term, lemma 3.1 and the Hölder inequality for the expectation value imply

$$\mathbb{E} \left[ \|1_{I_n} \gamma_{\Psi}^{(1)} 1_{I_m}\|_2^{\frac{2}{5}} \right] \leq 2^{\frac{1}{5}} \mathbb{E}[N(I_n)]^{\frac{1}{5}} \left( \sum_{k=1}^M \mathbb{E}[Q_L(k, m; J)] \right)^{\frac{2}{5}}. \tag{3.25}$$

The first factor is uniformly bounded according to (3.23). The last factor is bounded using the localization assumption (ECL):

$$\sum_{k=1}^M \mathbb{E}[Q_L(k, m; J)] \leq C \sum_{k=1}^M \exp \left( -\frac{\text{dist}(k, m)^{\xi}}{\ell^{\xi}} \right) \leq C' \exp \left( -\frac{\text{dist}(M, m)^{\xi}}{\ell^{\xi}} \right). \tag{3.26}$$

Since  $\text{dist}(M, m) \geq \frac{1}{2} \text{dist}(m, n)$  this implies the claim for the first term in the decomposition (3.6). The second term is treated similarly. For the third term, we employ lemma 3.2 and Hölder’s inequality for the expectation value to conclude:

$$\begin{aligned} &\mathbb{E} \left[ \|1_{I_n} \gamma_{\Psi}^{(3)} 1_{I_m}\|_2^{\frac{2}{5}} \right] \\ &\leq e^{1/5} 2^{\frac{3}{5}} \mathbb{E}[N(I_n)]^{\frac{1}{5}} \mathbb{E}[N(I_m)]^{\frac{1}{5}} \mathbb{E} \left[ \left( \sum'_{(k,l)} \sqrt{N(I_k)} Q_L(k, l; J) \right)^{2/3} \right]^{3/5} \\ &\leq e^{1/5} 2^{\frac{3}{5}} \mathbb{E}[N(I_n)]^{\frac{1}{5}} \mathbb{E}[N(I_m)]^{\frac{1}{5}} \sum'_{(k,l)} \mathbb{E}[N(I_k)]^{\frac{1}{5}} \mathbb{E}[Q_L(k, l; J)]^{\frac{2}{5}}. \end{aligned} \tag{3.27}$$

The terms involving the local particle number are uniformly bounded. The last term is again bounded using the localization assumption (ECL). In fact

$$\sum'_{(k,l)} \exp\left(-\frac{2\text{dist}(k, l)^\xi}{5\ell^\xi}\right) \leq C \exp\left(-\frac{2 \min\{\text{dist}(M, m)^\xi, \text{dist}(M, n)\}^\xi}{5\ell^\xi}\right). \tag{3.28}$$

Since the distance of the midpoint  $M$  to either  $n$  and  $m$  is at most  $\frac{1}{2}(1 + \text{dist}(n, m))$ , this completes the proof in case  $\sigma = \frac{2}{5}$ .

The general case follows with the help of interpolation from the bounds (3.21)–(3.23): For  $2/5 < \sigma < p$ ,

$$\begin{aligned} \mathbb{E}[\|1_{I_n} \gamma_\Psi 1_{I_m}\|_2^\sigma] &\leq \mathbb{E}\left[\|1_{I_n} \gamma_\Psi 1_{I_m}\|_2^{\frac{2(p-\sigma)}{5p-2}} N(I_n)^{\frac{p}{2} \frac{5\sigma-2}{5p-2}} N(I_m)^{\frac{p}{2} \frac{5\sigma-2}{5p-2}}\right] \\ &\leq \mathbb{E}\left[\|1_{I_n} \gamma_\Psi 1_{I_m}\|_2^{\frac{2}{5}}\right]^{\frac{5(p-\sigma)}{5p-2}} \mathbb{E}[N(I_n)^p]^{\frac{5\sigma-2}{2(5p-2)}} \mathbb{E}[N(I_m)^p]^{\frac{5\sigma-2}{2(5p-2)}}, \end{aligned} \tag{3.29}$$

where the second step is Hölder’s inequality for the expectation. Since the last two factors are uniformly bounded for  $p = 1$  we arrive at the claim.  $\square$

A proof of absence of BEC then proceeds as follows:

**Proof of corollary 2.3.** Assumption (2.11) implies that  $\sup_{L,n} \mathbb{E}[N(I_n)^p] < \infty$ . From the interpolation bound in (3.29) and (2.10) we conclude that for any  $2/5 < \sigma < p$

$$\mathbb{E}[\|1_{I_n} \gamma_\Psi 1_{I_m}\|_2^\sigma] \leq C \exp\left(-\frac{2(p-\sigma) \text{dist}(n, m)^\xi}{5p-2 (2\ell)^\xi}\right) \tag{3.30}$$

with some  $C < \infty$  that is independent of  $L, n$  and  $m$ . Since  $\|\gamma_\Psi\| \leq \max_n \sum_m \|1_{I_n} \gamma_\Psi 1_{I_m}\|$ , we can bound

$$\mathbb{E}[\|\gamma_\Psi\|^\sigma] \leq \sum_n \mathbb{E}\left[\left(\sum_m \|1_{I_n} \gamma_\Psi 1_{I_m}\|\right)^\sigma\right] \leq \sum_n \left(\sum_m \mathbb{E}[\|1_{I_n} \gamma_\Psi 1_{I_m}\|^\sigma]\right)^{1/\sigma}, \tag{3.31}$$

where in the last step we used Minkowski’s inequality for the expectation. Since the operator norm is bounded by the Hilbert–Schmidt norm, (3.30) implies that the sum over  $m$  in (3.31) is bounded, independently of  $n$ . This shows  $\mathbb{E}[\|\gamma_\Psi\|^\sigma] \leq CL$  with some  $C < \infty$  that is independent of  $L$  and  $N$ . A Chebychev estimate then implies for any  $\varepsilon > 0$  and  $r > 0$

$$\mathbb{P}(\|\gamma_\Psi\| > \varepsilon L^r) \leq \frac{\mathbb{E}[\|\gamma_\Psi\|^\sigma]}{\varepsilon^\sigma L^{r\sigma}} \leq \frac{C}{\varepsilon^\sigma} L^{1-\sigma r}, \tag{3.32}$$

where  $\mathbb{P}$  stands for the probability of an event. If we choose  $r > 2/\sigma$ , then  $1 - \sigma r < -1$ , and the right side of (3.32) is summable in  $L$ . The Borel–Cantelli lemma thus yields the claimed almost-sure convergence.  $\square$

### 4. Proof of the absence of superfluidity

For a proof of theorem 2.4, let  $H_L(\theta)$  denote the self-adjoint operator which acts as (1.2) on functions with twisted b.c.s,  $\psi(L) = e^{i\theta}\psi(0)$  and  $\psi'(L) = e^{i\theta}\psi'(0)$ . Let  $\{E_{j,L}(\theta)\}$  denote its eigenvalues, ordered increasingly with  $j$ , i.e.,  $E_{j,L}(\theta) \leq E_{j+1,L}(\theta)$ . With

$$\theta_\mu := \theta + \frac{\pi}{2}(1 + (-1)^{N_\mu}) \tag{4.1}$$

we have

$$E_L(N_\mu, \theta) = \sum_{j=1}^{N_\mu} E_{j,L}(\theta_\mu). \tag{4.2}$$

We claim that  $E_{N_\mu,L}(\theta_\mu) \leq \mu$ . This follows from the fact that, by construction  $E_{N_\mu,L}^\pm \leq \mu$ , and  $E_{N_\mu,L}(\theta_\mu) \leq \max\{E_{N_\mu,L}^+, E_{N_\mu,L}^-\}$ . Hence we can invoke the variational principle in the form

$$E_L(N_\mu, \theta) = \mu N_\mu + \inf\{\text{Tr}[H_L(\theta_\mu) - \mu]\gamma \mid 0 \leq \gamma \leq 1, \text{Tr} \gamma \leq N_\mu\}. \tag{4.3}$$

We emphasize that it is possible here to relax the condition  $\text{Tr} \gamma = N_\mu$  to  $\text{Tr} \gamma \leq N_\mu$  exactly because  $E_{N_\mu,L}(\theta_\mu) \leq \mu$ . This turns out to be convenient in the following.

To obtain an upper bound on  $E_L(N_\mu, \theta)$ , and hence on  $\rho_s$ , we choose as a trial density matrix in (4.3)

$$\gamma = \frac{\tilde{\gamma}}{\max\{\|\tilde{\gamma}\|, 1\}}, \quad \text{with} \quad \tilde{\gamma} := \sum_{j: E_{j,L}^\mu \leq \mu} e^{i\psi_{j,L}} |\varphi_{j,L}^\mu\rangle \langle \varphi_{j,L}^\mu| e^{-i\psi_{j,L}}. \tag{4.4}$$

Here,  $\{\varphi_{j,L}^\mu\}$  abbreviates an orthonormal eigenbasis of  $H_L^\mu$  and  $\{E_{j,L}^\mu\}$  are the corresponding eigenvalues. Note that  $\text{Tr} \tilde{\gamma} = N_\mu$ , as remarked in (1.9), hence  $0 \leq \gamma \leq 1$  and  $\text{Tr} \gamma \leq N_\mu$ .

The trial phase functions  $\psi_{j,L} : [0, L] \rightarrow \mathbb{R}$  will be chosen continuous, increasing (and piecewise differentiable) such that  $\psi_{j,L}(0) = 0$  and  $\psi_{j,L}(L) = \theta$ . We pick them depending on an additional variational parameter  $\delta > 0$  to be chosen later. More specifically, we set

$$I_j(\delta) := \bigcup_{k: \Phi_{j,L}^\mu(k) \leq \delta} I_k, \tag{4.5}$$

and choose

$$\tilde{I}_j(\delta) \subseteq I_j(\delta) \tag{4.6}$$

to be the largest connected subset of  $I_j(\delta)$ . We then simply take  $\psi_{j,L}$  to increase linearly on  $\tilde{I}_j(\delta)$  with slope  $\theta/|\tilde{I}_j(\delta)|$ , and constant otherwise. Note that  $\tilde{I}_j(\delta)$  is certainly non-empty for  $\delta > L^{-1/2}$ , since  $1 \geq \sum_{n: \Phi_{j,L}^\mu(n) > \delta} \Phi_{j,L}^\mu(n)^2 \geq (L - |I_j(\delta)|) \delta^2$  which implies  $|I_j(\delta)| \geq L - \delta^{-2}$ . The localization assumption (SULE) may be used for a stronger estimate. Namely, (2.6) implies that  $\Phi_{j,L}^\mu(k) \leq \delta$  for any  $k \in \{1, \dots, L\}$  with  $\text{dist}(k, \gamma_{j,L}^\mu)^\xi \geq \ell^\xi \ln(A_L L^{3/2}/\delta)$ . Choosing

$$\delta := A_L L^{3/2} \exp\left(-\frac{L^\xi}{(4\ell)^\xi}\right), \tag{4.7}$$

we see that  $\Phi_{j,L}^\mu(k) \leq \delta$  whenever  $\text{dist}(k, \gamma_{j,L}^\mu) \geq L/4$ , and thus  $I_j(\delta)$  contains an interval of length at least  $(L - 4)/4$ . Therefore

$$|\tilde{I}_j(\delta)| \geq \frac{L - 4}{4}. \tag{4.8}$$

A straightforward computation shows that

$$\begin{aligned} & \text{Tr}[H_L(\theta_\mu) - \mu]\gamma \\ &= -\frac{\text{Tr}[H_L^\mu - \mu]_-}{\max\{\|\tilde{\gamma}\|, 1\}} + \frac{1}{\max\{\|\tilde{\gamma}\|, 1\}} \sum_{j: E_{j,L}^\mu \leq \mu} \int_0^L |\varphi_{j,L}^\mu(x)|^2 \left| \frac{d}{dx} \psi_{j,L}(x) \right|^2 dx \\ &\leq -\frac{\text{Tr}[H_L^\mu - \mu]_-}{\max\{\|\tilde{\gamma}\|, 1\}} + \theta^2 \sum_{j: E_{j,L}^\mu \leq \mu} \frac{1}{|\tilde{I}_j(\delta)|^2} \sum_{n: I_n \subset \tilde{I}_j(\delta)} \Phi_{j,L}^\mu(n)^2 \\ &\leq -\frac{\text{Tr}[H_L^\mu - \mu]_-}{\max\{\|\tilde{\gamma}\|, 1\}} + \theta^2 \delta^2 \sum_{j: E_{j,L}^\mu \leq \mu} \frac{1}{|\tilde{I}_j(\delta)|}. \end{aligned} \tag{4.9}$$

To estimate the norm of  $\tilde{\gamma}$ , we note that  $\tilde{\gamma}$  is unitarily equivalent to the  $N_\mu \times N_\mu$  matrix with matrix elements  $\langle \varphi_{j,L}^\mu e^{i\psi_{j,L}}, e^{i\psi_{k,L}} \varphi_{k,L}^\mu \rangle$ . In particular

$$\|\tilde{\gamma}\| \leq \max_j \sum_k |\langle \varphi_{j,L}^\mu e^{i\psi_{j,L}}, e^{i\psi_{k,L}} \varphi_{k,L}^\mu \rangle|. \tag{4.10}$$

For  $j \neq k$ , we have

$$\begin{aligned} & \int_{\tilde{I}_j(\delta)^c \cap \tilde{I}_k(\delta)^c} \overline{\varphi_{j,L}^\mu(z)} e^{-i(\psi_{j,L}(z) - \psi_{k,L}(z))} \varphi_{k,L}^\mu(z) dz \\ &= e^{i\alpha} \int_{\tilde{I}_j(\delta)^c \cap \tilde{I}_k(\delta)^c} \overline{\varphi_{j,L}^\mu(z)} \varphi_{k,L}^\mu(z) dz \\ &= -e^{i\alpha} \int_{\tilde{I}_j(\delta) \cup \tilde{I}_k(\delta)} \overline{\varphi_{j,L}^\mu(z)} \varphi_{k,L}^\mu(z) dz \end{aligned} \tag{4.11}$$

since  $\psi_{k,L}(z) - \psi_{j,L}(z) = \alpha$  is a constant on  $\tilde{I}_j(\delta)^c \cap \tilde{I}_k(\delta)^c$ . Hence

$$\begin{aligned} & |\langle \varphi_{j,L}^\mu e^{i\psi_{j,L}}, e^{i\psi_{k,L}} \varphi_{k,L}^\mu \rangle| \\ &\leq \int_{\tilde{I}_j(\delta) \cup \tilde{I}_k(\delta)} |e^{-i(\psi_{j,L}(z) - \psi_{k,L}(z))} - e^{i\alpha}| |\varphi_{j,L}^\mu(z) \varphi_{k,L}^\mu(z)| dz \\ &\leq 2 \max_{|\beta| \leq |\theta|} |1 - e^{i\beta}| \delta L \leq 2|\theta| \delta L \end{aligned} \tag{4.12}$$

for  $j \neq k$  and  $|\theta| \leq \pi$ . In combination with (4.10), this implies

$$\|\tilde{\gamma}\| \leq 1 + 2|\theta| \delta L N_\mu. \tag{4.13}$$

Inserting the bounds (4.8) and (4.13) in (4.9), and using (4.3) and (1.9), we hence conclude that for any  $L \geq 5$

$$X_L := L \frac{E_L(N_\mu, \theta) - E_L(N_\mu, 0)}{\theta^2} \leq \frac{2\delta L^2}{|\theta|} \text{Tr}[H_L^\sharp - \mu]_- N_\mu + \frac{4L\delta^2}{L-4} N_\mu, \tag{4.14}$$

with  $\delta$  given in (4.7). Our goal is to show that  $\lim_{L \rightarrow \infty} X_L = 0$  almost surely. Note that  $\delta$  is random, but the coefficient  $A_L$  in (4.7) is uniformly bounded in expectation, according to the assumption (2.5). Moreover, we have the following rough but uniform bounds:

**Lemma 4.1.** *Under assumption (1.3) one has, irrespective of b.c.s*

$$\sup_{L \in \mathbb{N}} \frac{\mathbb{E}[N_\mu]}{L} < \infty \quad \text{and} \quad \sup_{L \in \mathbb{N}} \frac{\mathbb{E}[(\text{Tr}[H_L^\sharp - \mu]_-)^{1/3}]}{L} < \infty. \tag{4.15}$$

**Proof.** With  $h_n$  denoting the restriction of  $H_L$  to  $I_n$ , with Neumann b.c.s, it is well known [47, section XIII.15] that

$$N_\mu \leq \sum_{n=1}^L \text{Tr} \mathbf{1}_{(-\infty, \mu]}(h_n). \tag{4.16}$$

A simple calculation based on the Birman–Schwinger principle (see, e.g., [51, chapter 7]) shows that  $\inf \text{spec } h_n$  can be bounded in terms of  $\int_{I_n} |V_-|$ , which has a finite expectation according to our assumption (1.3). In fact, one has

$$\inf \text{spec } h_n \geq -f^{-1}\left(\int_{I_n} |V_-|\right), \quad f(t) = \sqrt{t} \tanh \sqrt{t}, \tag{4.17}$$

where  $V_-$  denotes the negative part of  $V$ . Keeping half of the kinetic energy, one also obtains

$$h_n \geq -\frac{1}{2} \Delta_{I_n} - \frac{1}{2} f^{-1}\left(2 \int_{I_n} |V_-|\right), \tag{4.18}$$

and hence

$$\text{Tr} \mathbf{1}_{(-\infty, \mu]}(h_n) \leq \text{Tr} \mathbf{1}_{[0, 2\mu + f^{-1}(2 \int_{I_n} |V_-|)]}(-\Delta_{I_n}). \tag{4.19}$$

It is easy to see that  $\text{Tr} \mathbf{1}_{[0, \nu]}(-\Delta_{I_n})$  grows like  $\nu^{1/2}$  for large  $\nu$ , and  $f^{-1}(t)$  grows like  $t^2$  for large  $t$ . Hence (4.19) is bounded by a constant times  $1 + \int_{I_n} |V_-|$ , which implies, in combination with (4.16), the first bound in (4.15).

To obtain the second, we use, similarly to (4.16), that

$$\text{Tr}[H_L^\sharp - \mu]_- \leq -\sum_{n=1}^L \text{Tr}(h_n - \mu) \mathbf{1}_{(-\infty, \mu]}(h_n). \tag{4.20}$$

In combination with (4.17) this implies

$$\text{Tr}[H_L^\sharp - \mu]_- \leq \sum_{n=1}^L \left( \mu + f^{-1}\left(\int_{I_n} |V_-|\right) \right) \text{Tr} \mathbf{1}_{(-\infty, \mu]}(h_n). \tag{4.21}$$

In particular,  $(\text{Tr}[H_L^\sharp - \mu]_-)^{1/3}$  is bounded by a constant times  $\sum_n (1 + \int_{I_n} |V_-|)$ , which implies the desired result (4.15). □

Let us denote the right side of (4.14) by  $Y_L$ . From lemma 4.1, (2.5) and Hölder’s inequality for the expectation value, it follows that

$$\mathbb{E}[Y_L^{1/5}] \leq C |\theta|^{-1/5} L^{3/2} \exp\left(-\frac{1}{5} \frac{L^\xi}{(4\ell)^\xi}\right) \tag{4.22}$$

for  $L \geq 5$  and some constant  $C > 0$  independent of  $L$ . The Chebychev inequality yields, for any  $\varepsilon > 0$ ,

$$\mathbb{P}(X_L \geq \varepsilon) \leq \mathbb{P}(Y_L \geq \varepsilon) \leq \varepsilon^{-1/5} \mathbb{E}[Y_L^{1/5}]. \tag{4.23}$$

From (4.22), the right side is seen to be summable in  $L$ . The proof of theorem 2.4 is thus concluded with the help of the Borel–Cantelli lemma, which ensures that the probability that  $X_L \geq \varepsilon$  happens for infinitely many  $L \in \mathbb{N}$  is zero. □

### Acknowledgments

Part of this work was carried out at the Erwin Schrödinger Institute (ESI) in Vienna, Austria, during the programme ‘Quantum many-body systems, random matrices, and disorder’. We thank the ESI for hospitality and financial support, and W Zwerger for useful comments. RS acknowledges financial support by the Austrian

Science Fund (FWF), project No. P 27533-N27, and SW by the Deutsche Forschungsgemeinschaft (WA 1699/2-1).

## Appendix. Proof of dynamical properties of the density

**Proof of proposition 2.1.** By assumption on the range of the initial state, we have  $\Gamma_{\phi_t} = U_t^* \Gamma_{\phi} U_t$  with  $U_t = e^{itH_t^N} P_J(H_t^{\#N})$ . Consequently

$$\begin{aligned} \left| \int_I \varrho_t(x) dx - \int_I \varrho_0(x) dx \right| &= |\text{Tr } 1_I U_t^* \Gamma_{\phi} U_t - \text{Tr } 1_I \Gamma_{\phi}| \\ &= |\text{Tr } 1_I U_t^* 1_{I^c} \Gamma_{\phi} U_t - \text{Tr } 1_{I^c} U_t^* 1_I \Gamma_{\phi} U_t| \\ &\leq \|1_I U_t^* 1_{I^c}\|_1 + \|1_{I^c} U_t^* 1_I\|_1, \end{aligned} \quad (\text{A.1})$$

where the inequality follows from  $\|\Gamma_{\phi} U_t\| \leq \|\Gamma_{\phi}\| \leq 1$ . The first bound (2.8) is then a consequence of

$$\mathbb{E}[\sup_{t \in \mathbb{R}} \|1_{K^c} U_t^* 1_I\|_1] \leq \sum_{\substack{I_n \cap I \neq \emptyset \\ I_m \cap K^c \neq \emptyset}} \mathbb{E}[\sup_{t \in \mathbb{R}} \|1_{I_n} U_t^* 1_{I_m}\|_1], \quad (\text{A.2})$$

valid for all  $I \subseteq K$ . In case  $I = K$ , the right side is bounded by a constant on account of (2.4); this concludes the proof of the first assertion.

The proof of the second assertion (2.9) proceeds similarly. We estimate

$$\begin{aligned} \int_I \varrho_t(x) dx &= \text{Tr } 1_I U_t^* 1_{K^c} \Gamma_{\phi} 1_K U_t + \text{Tr } 1_I U_t^* 1_K \Gamma_{\phi} 1_{K^c} U_t + \text{Tr } 1_I U_t^* 1_{K^c} \Gamma_{\phi} U_t \\ &\leq \|1_K \Gamma_{\phi} 1_{K^c}\|_1 + \|1_{K^c} U_t 1_I\|_1 + \|1_I U_t^* 1_{K^c}\|_1 = \int_K \varrho_0(x) dx + 2\|1_{K^c} U_t 1_I\|_1. \end{aligned} \quad (\text{A.3})$$

The proof is completed using (A.2) and (2.4).  $\square$

## References

- [1] Abdul-Rahman H and Stolz G 2015 A uniform area law for the entanglement of eigenstates in the disordered XY chain *J. Math. Phys.* **56** 121901
- [2] Abdul-Rahman H, Nachtergaele B, Sims R and Stolz G 2015 Entanglement dynamics of disordered quantum XY chains arXiv:1510.00262
- [3] Aizenman M and Molchanov S 1993 Localization at large disorder and at extreme energies: an elementary derivation *Commun. Math. Phys.* **157** 245–78
- [4] Aizenman M, Elgart A, Naboko S, Schenker J and Stolz G 2006 Moment analysis for localization in random Schrödinger operators *Invent. Math.* **163** 343–413
- [5] Aizenman M and Warzel S 2016 *Random Operators: Disorder Effects on Quantum Spectra and Dynamics* (Providence, RI: American Mathematical Society) GSM 168
- [6] Altman E and Vosk R 2015 Universal dynamics and renormalization in many-body-localized systems *Annu. Rev. Condens. Matter Phys.* **6** 383–409
- [7] Barth M and Zwerger W 2011 Tan relations in one dimension *Ann. Phys., NY* **326** 2544–65
- [8] Bishop M and Wehr J 2013 Ground state energy of mean-field model of interacting bosons in Bernoulli potential *J. Math. Phys.* **54** 081902
- [9] Cazalilla M A, Citro R, Giamarchi T, Orignac E and Rigol M 2011 One dimensional bosons: from condensed matter systems to ultracold gases *Rev. Mod. Phys.* **83** 1405–66
- [10] del Rio R, Jitomirskaya S, Last Y and Simon B 1995 What is localization? *Phys. Rev. Lett.* **75** 117–9
- [11] De Martino A, Thorwart M, Egger R and Graham R 2005 Exact results for one-dimensional disordered bosons with strong repulsion *Phys. Rev. Lett.* **94** 060402
- [12] Eastham M S P 1973 *The Spectral Theory of Periodic Differential Equations* (Edinburgh: Scottish Academic)
- [13] Fischer W, Leschke H and Müller P 2000 Spectral localization by Gaussian random potentials in multi-dimensional continuous space *J. Stat. Phys.* **101** 935–85
- [14] Fröhlich J and Spencer T 1983 Absence of diffusion in the Anderson tight binding model for large disorder or low energy *Commun. Math. Phys.* **88** 151–84
- [15] Fisher M E, Barber M N and Jasnow D 1973 Helicity modulus, superfluidity, and scaling in isotropic systems *Phys. Rev. B* **8** 1111–24
- [16] Fisher M P A, Weichman P B, Grinstein G and Fisher D S 1989 Boson localization and the superfluid-insulator transition *Phys. Rev. B* **40** 546–70
- [17] Germinet F and Klein A 2001 Bootstrap multiscale analysis and localization in random media *Commun. Math. Phys.* **222** 415–48
- [18] Germinet F and Klein A 2006 New characterizations of the region of complete localization for random Schrödinger operators *J. Stat. Phys.* **122** 73–94
- [19] Giamarchi T and Schulz H J 1988 Anderson localization and interactions in one-dimensional metals *Phys. Rev. B* **37** 325–40
- [20] Girardeau M 1960 Relationship between systems of impenetrable bosons and fermions in one dimension *J. Math. Phys.* **1** 516–23
- [21] Goldscheid I Ya, Molchanov S and Pastur L 1977 A pure point spectrum of the stochastic one-dimensional Schrödinger operator *Funct. Anal. Appl.* **11** 1–8
- [22] Hamza E, Sims R and Stolz G 2010 A note on fractional moments for the one-dimensional continuum Anderson model *J. Math. Anal. Appl.* **365** 435–46
- [23] Hamza E, Sims R and Stolz G 2012 Dynamical localization in disordered quantum spin systems *Commun. Math. Phys.* **315** 215–39

- [24] Imbrie J 2014 On many-body localization for quantum spin chains arXiv:1403.7837
- [25] Kac M and Luttinger J M 1973 Bose–Einstein condensation in the presence of impurities *J. Math. Phys.* **14** 1626–8
- [26] Kac M and Luttinger J M 1974 Bose–Einstein condensation in the presence of impurities: II. *J. Math. Phys.* **15** 183–6
- [27] Kinoshita T, Wenger T and Weiss D S 2004 Observation of a one-dimensional Tonks–Girardeau gas *Science* **305** 1125–8
- [28] Klein A and Perez J F 1990 Localization in the ground-state of the one dimensional XY model with a random transverse field *Commun. Math. Phys.* **128** 99–108
- [29] Klopp F and Veniaminov N 2014 Interacting electrons in a random medium: a simple one-dimensional model (arXiv:1408.5839)
- [30] Könenberg M, Moser T, Seiringer R and Yngvason J 2015 Superfluid behavior of a Bose–Einstein condensate in a random potential *New J. Phys.* **17** 013022
- [31] Lenard A 1964 Momentum distribution in the ground state of the one-dimensional system of impenetrable bosons *J. Math. Phys.* **5** 930–43
- [32] Lenoble O, Pastur L A and Zagrebnov V A 2004 Bose–Einstein condensation in random potentials *C. R. Phys.* **5** 129–42
- [33] Lenoble O and Zagrebnov V A 2007 Bose–Einstein condensation in the Luttinger–Sy model *Markov Process. Relat. Fields* **13** 441–68
- [34] Lieb E H and Liniger W 1962 Exact analysis of an interacting Bose gas: I. The general solution and the ground state *Phys. Rev.* **130** 1605–16
- [35] Lieb E H and Loss M 2001 *Analysis* 2nd edn (Providence, RI: American Mathematical Society)
- [36] Lieb E H, Seiringer R, Solovej J P and Yngvason J 2005 *The Mathematics of the Bose Gas and its Condensation (Oberwolfach Seminar Series vol 34)* (Basel: Birkhäuser)
- [37] Lifshits I, Gredeskul S and Pastur L 1988 *Introduction to the Theory of Disordered Systems* (New York: Wiley)
- [38] Lugan P, Clément D, Bouyer P, Aspect A, Lewenstein M and Sanchez-Palencia L 2007 Ultracold Bose gases in 1D disorder: from Lifshits glass to Bose–Einstein condensate *Phys. Rev. Lett.* **98** 170403
- [39] Luttinger J M and Sy H K 1973 Bose–Einstein condensation in a one-dimensional model with random impurities *Phys. Rev. A* **7** 712–20
- [40] Nandkishore R and Huse D A 2015 Many-body localization and thermalization in quantum statistical mechanics *Annu. Rev. Condens. Matter Phys.* **6** 15–38
- [41] Mastropietro V 2015 Localization of interacting fermions in the Aubry–André model *Phys. Rev. Lett.* **115** 180401
- [42] Paredes B, Widera A, Murg V, Mandel O, Fölling S, Cirac I, Shlyapnikov G V, Hänsch T W and Bloch I 2004 Tonks–Girardeau gas of ultracold atoms in an optical lattice *Nature* **429** 277–81
- [43] Pastur L and Figotin A 1992 *Spectra of Random and Almost-Periodic Operators* (Berlin: Springer)
- [44] Penrose O and Onsager L 1956 Bose–Einstein condensation and liquid helium *Phys. Rev.* **104** 576–84
- [45] Radić J, Bačić V, Jukić D, Segev M and Buljan I H 2010 Anderson localization of a Tonks–Girardeau gas in potentials with controlled disorder *Phys. Rev. A* **81** 063639
- [46] Rapsch S, Schollwöck U and Zwerger W 1999 Density matrix renormalization group for disordered bosons in one dimension *Eur. Phys. Lett.* **46** 559
- [47] Reed M and Simon B 1978 *Analysis of Operators* (New York: Academic)
- [48] Rigol M and Muramatsu A 2004 Emergence of quasi-condensates of hard-core bosons at finite momentum *Phys. Rev. Lett.* **93** 230404
- [49] Sanchez-Palencia L and Lewenstein M 2010 Disordered quantum gases under control *Nat. Phys.* **6** 87–95
- [50] Seiringer R, Yngvason J and Zagrebnov V A 2012 Disordered Bose–Einstein condensates with interaction in one dimension *J. Stat. Mech.* **P11007**
- [51] Simon B 2005 *Trace Ideals and Their Applications* 2nd edn (Providence, RI: American Mathematical Society)
- [52] Sims R and Stolz G 2015 Many-body localization: concepts and simple models *Markov Process. Relat. Fields* **21** 791–822
- [53] Sims R and Warzel S 2015 Decay of determinantal and pfaffian correlation functionals in one-dimensional lattices *Commun. Math. Phys.* (arXiv:1509.00450)
- [54] Stasinska J, Massingnan P, Bishop M, Wehr J, Sanpera A and Lewenstein M 2012 The glass to superfluid transition in dirty bosons on a lattice *New J. Phys.* **14** 043043
- [55] Ueki N 2004 Wegner estimates and localization for Gaussian random potentials *Publ. Res. Inst. Math. Sci.* **40** 29–90
- [56] Yukalov V I and Girardeau M D 2005 Fermi–Bose mapping for one-dimensional Bose gases *Laser Phys. Lett.* **2** 375–82