



Multiple Genome Sequences of Exopolysaccharide-Producing, Brewery-Associated *Lactobacillus brevis* Strains

Marion E. Fraunhofer, Andreas J. Geissler, Frank Jakob, Rudi F. Vogel

Technische Universität München, Lehrstuhl für Technische Mikrobiologie, Freising, Germany

ABSTRACT *Lactobacillus brevis* represents one of the most relevant beer-spoiling bacteria. Besides strains causing turbidity and off flavors upon growth and metabolite formation, this species also comprises strains that produce exopolysaccharides (EPSs), which increase the viscosity of beer. Here, we report the complete genome sequences of three EPS-producing, brewery-associated *L. brevis* strains.

Beer represents a microbiologically stable beverage, as it combines a low pH and nutrient availability with the presence of hop compounds, ethanol, carbon dioxide, and anaerobicity (1). Nevertheless, certain lactobacilli are still able to grow in beer and spoil it. The resulting spoilage manifests as acidity, turbidity, off flavor, or increased viscosity. The latter is attributed to exopolysaccharides (EPSs) derived from members of the family *Lactobacillaceae* (2). To gain insights into this EPS synthesis, we sequenced the complete genomes of three brewery-associated *Lactobacillus brevis* strains.

L. brevis TMW 1.2108 and *L. brevis* TMW 1.2111 were isolated from wheat beer, and *L. brevis* TMW 1.2113 was isolated from a brewery surface. The Genomic-tip 100/G kit (Qiagen, Hilden, Germany) was used to isolate high-molecular-weight DNA from MRS liquid cultures. Single-molecule real-time (SMRT) sequencing (PacBio RS II) was performed at GATC (Konstanz, Germany) (3). For each of the three strains, a single library was prepared by selecting an insert size of 8 to 12 kb, resulting in at least 200 Mb of raw data from one SMRT cell (1 × 120-min movies), applying P4-C2 chemistry. The generated sequences were assembled with SMRT Analysis version 2.2.0.p2 using the Hierarchical Genome Assembly Process (HGAP) version 3 (4). The genome was completed by manual curation according to PacBio instructions and annotated using the NCBI Prokaryotic Genome Annotation Pipeline (PGAP) and the Rapid Annotations using Subsystems Technology (RAST) server (5–7).

Table 1 summarizes the characteristics, sequencing statistics, genome information, and accession numbers for each strain. The chromosome sizes range from 2.54 to 2.57 Mb, with G+C contents of 45.8% to 45.9%. The strains harbor 4 to 8 plasmids with G+C contents between 41.3% and 41.9% and sizes between 130.5 and 352.0 kb. The chromosomes encode five complete rRNA operons and 66 to 69 tRNAs.

All strains possess a plasmid-encoded glycosyltransferase-2 (*gtf-2*), which has been described as a key enzyme for EPS synthesis of slimy, wine-spoiling members of the family *Lactobacillaceae* (8–10). Comparison of the *gtf-2* gene between the wine spoilers *Pediococcus parvulus* IOEB8801 (GenBank accession no. AF196967), *P. damnosus* 2.6 (GenBank accession no. AY999683), and *L. diolivorans* G77 (GenBank accession no. AY999684) and the beer spoilers reported here reveals sequence identities of 99%. This shows the *gtf-2* gene to be species-independent and highly conserved and might indicate a common origin.

The availability of these *L. brevis* genome sequences will allow a better understanding of EPS synthesis and its contribution to the spoilage of beer.

Received 4 May 2017 Accepted 9 May 2017 Published 29 June 2017

Citation Fraunhofer ME, Geissler AJ, Jakob F, Vogel RF. 2017. Multiple genome sequences of exopolysaccharide-producing, brewery-associated *Lactobacillus brevis* strains. *Genome Announc* 5:e00585-17. <https://doi.org/10.1128/genomeA.00585-17>.

Copyright © 2017 Fraunhofer et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Frank Jakob, frank.jakob@wzw.tum.de.

TABLE 1 Strain characteristics, sequencing statistics, genome information, and accession numbers

Strain	Source	BioSample no. ^a	Accession no. ^b	Coverage (×) ^c	Size (Mb)	No. of contigs ^d	G+C content (%)	No. of CDSs ^e
<i>L. brevis</i> TMW 1.2108	Beer	SAMN04517635	CP019734 to CP019742	148	2.92	9	45.29	2,582
<i>L. brevis</i> TMW 1.2111	Beer	SAMN04517636	CP019743 to CP019749	341	2.88	7	45.32	2,440
<i>L. brevis</i> TMW 1.2113	Brewery surface	SAMN04517634	CP019750 to CP019754	364	2.67	8	45.74	2,357

^aAll BioSamples are part of BioProject no. [PRJNA313253](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA313253).

^bAccession numbers are listed for all contigs of each whole genome (as a range).

^cAverage coverage of HGAP assembly.

^dIn the chromosomes plus plasmids and partial plasmids.

^eCDSs, number of coding sequences (total) based on NCBI PGAP.

Accession number(s). The three complete *L. brevis* genomes have been deposited in DDBJ/EMBL/GenBank under the accession numbers given in Table 1.

ACKNOWLEDGMENTS

This work was supported by the German Research Foundation (DFG) and the Technische Universität München within the funding program Open Access Publishing. Parts of this work were funded by the German Ministry of Economics and Technology (via AiF) and the Wifoe (Wissenschaftsförderung der Deutschen Brauwirtschaft e.V., Berlin), project AiF 18194N. None of the funding sources had any influence on the study design, the collection, analysis and interpretation of data, the writing of the report, and the decision to submit the article for publication.

REFERENCES

- Suzuki K. 2011. 125th anniversary review: microbiological instability of beer caused by spoilage bacteria. *J Inst Brew* 117:131–155. <https://doi.org/10.1002/j.2050-0416.2011.tb00454.x>.
- Williamson DH. 1959. Studies on lactobacilli causing ropiness in beer. *J Appl Bacteriol* 22:11.
- Eid J, Fehr A, Gray J, Luong K, Lyle J, Otto G, Peluso P, Rank D, Baybayan P, Bettman B, Bibillo A, Bjornson K, Chaudhuri B, Christians F, Cicero R, Clark S, Dalal R, Dewinter A, Dixon J, Foquet M, Gaertner A, Hardenbol P, Heiner C, Hester K, Holden D, Kearns G, Kong X, Kuse R, Lacroix Y, Lin S, Lundquist P, Ma C, Marks P, Maxham M, Murphy D, Park I, Pham T, Phillips M, Roy J, Sebra R, Shen G, Sorenson J, Tomaney A, Travers K, Trulson M, Vieceli J, Wegener J, Wu D, Yang A, Zaccarin D. 2009. Real-time DNA sequencing from single polymerase molecules. *Science* 323:133–138. <https://doi.org/10.1126/science.1162986>.
- Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Non-hybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <https://doi.org/10.1038/nmeth.2474>.
- Angiuoli SV, Gussman A, Klimke W, Cochrane G, Field D, Garrity G, Kodira CD, Kyrpides N, Madupu R, Markowitz V, Tatusova T, Thomson N, White O. 2008. Toward an online repository of standard operating procedures (SOPs) for (meta)genomic annotation. *Omics* 12:137–141. <https://doi.org/10.1089/omi.2008.0017>.
- Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: rapid annotations using subsystems technology. *BMC Genomics* 9:75. <https://doi.org/10.1186/1471-2164-9-75>.
- Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the rapid annotation of microbial genomes using subsystems technology (RAST). *Nucleic Acids Res* 42:D206–D214. <https://doi.org/10.1093/nar/gkt1226>.
- Dols-Lafargue M, Lee HY, Le Marrec C, Heyraud A, Chambat G, Lonvaud-Funel A. 2008. Characterization of *gtf*, a glucosyltransferase gene in the genomes of *Pediococcus parvulus* and *Oenococcus oeni*, two bacterial species commonly found in wine. *Appl Environ Microbiol* 74:4079–4090. <https://doi.org/10.1128/AEM.00673-08>.
- Walling E, Gindreau E, Lonvaud-Funel A. 2005. A putative glucan synthase gene *dps* detected in exopolysaccharide-producing *Pediococcus damnosus* and *Oenococcus oeni* strains isolated from wine and cider. *Int J Food Microbiol* 98:53–62. <https://doi.org/10.1016/j.ijfoodmicro.2004.05.016>.
- Werning ML, Ibarburu I, Dueñas MT, Irastorza A, Navas J, López P. 2006. *Pediococcus parvulus* *gtf* gene encoding the GTF glucosyltransferase and its application for specific PCR detection of beta-D-glucan-producing bacteria in foods and beverages. *J Food Prot* 69:161–169. <https://doi.org/10.4315/0362-028X-69.1.161>.