

# Subpectoral Biceps Tenodesis: All-Suture Anchor Onlay Technique



Lucca Lacheta, M.D., Andreas B. Imhoff, M.D., Sebastian Siebenlist, M.D., and Bastian Scheiderer, M.D.

**Abstract:** Surgical management for biceps pathologies has advanced to provide stronger fixation and to be less invasive. The long head of the biceps tendon has been recognized as a common contributor to anterior shoulder pain and is often associated with other glenohumeral pathologies such as SLAP lesions, rotator cuff tears, and subacromial impingement. Both tenotomy and tenodesis have shown to be effective in ameliorating pain associated with the long head of the biceps tendon. However, decreased muscle function and cosmetic concerns are seen at higher rates after tenotomy compared with tenodesis. One option for the treatment of biceps tendon pathology includes mini-open subpectoral biceps tenodesis. Lower reoperation rates are observed after subpectoral biceps tenodesis than after suprapectoral biceps tenodesis, with thoughts that releasing the tendon from its sheath and the bicipital groove relieves the patient of most associated pain. The purpose of this Technical Note is to describe in detail our preferred operative technique for mini-open subpectoral biceps tenodesis using an onlay technique with all-suture anchor fixation.

**P**athologies of the long head of the biceps (LHB) tendon are a common contributor to anterior shoulder pain.<sup>1-6</sup> Biceps tendinopathy can arise from various causes, including chronic tendinitis, acute trauma, overuse, and degenerative changes. Patients present with pain during overhead movements and internal rotation, as well as pain-limited range of motion.<sup>6</sup>

Both tenotomy and tenodesis have shown to be effective in ameliorating pain associated with the LHB tendon. However, decreased muscle function and cosmetic defects are observed at higher rates after tenotomy compared with tenodesis.<sup>4,5,7-9</sup> In addition, lower reoperation rates are described after subpectoral fixation than after suprapectoral fixation, with thoughts that releasing the

tendon from its sheath and the bicipital groove relieves the patient of most associated pain.<sup>5,10</sup>

There is no clear consensus whether bone tunnel or cortical surface (onlay) healing confers better outcomes. Clinical studies comparing interference screw fixation (intramedullary) and suture anchor fixation (onlay) techniques for subpectoral biceps tenodesis found no significant differences in patient outcomes.<sup>10,11</sup> However, interference screw technique has been associated with various complications including humeral fractures at the drill hole, persistent pain, and bioabsorbable screw reactions.<sup>12-14</sup> Subpectoral biceps tenodesis using an onlay fixation technique is a reasonable alternative to mitigate these risks.

The purpose of this Technical Note is to describe in detail our preferred technique for mini-open subpectoral biceps tenodesis using an all-suture anchor onlay fixation technique. We believe that this technique is a viable option for the treatment of biceps tendinopathy with a potentially decreased risk of humeral fractures and its easy approach and anchor placement at the humeral shaft using curved drilling and insertion guides.

## Surgical Technique

### Patient Positioning

After induction of general anesthesia, the patient is placed in the beach-chair position. The operative extremity is prepared and draped in standard sterile

*From the Department of Orthopaedic Sports Medicine, Technical University of Munich, Munich, Germany.*

*The authors report the following potential conflicts of interest or sources of funding: L.L. receives nonfinancial support for a position at Steadman Hippo Research Institute supported by Arthrex. Full ICMJE author disclosure forms are available for this article online, as [supplementary material](#).*

*Received October 25, 2019; accepted January 11, 2020.*

*Address correspondence to Bastian Scheiderer, M.D., Department of Orthopaedic Sports Medicine, Technical University of Munich, Ismaninger Strasse 22, 81675 Munich, Germany. E-mail: [bastian.scheiderer@tum.de](mailto:bastian.scheiderer@tum.de)*

*© 2020 by the Arthroscopy Association of North America. Published by Elsevier. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).*

*2212-6287/191301*

*<https://doi.org/10.1016/j.eats.2020.01.019>*




**Fig 1.** The right arm is positioned in 90° of abduction and 90° of elbow flexion. A 3-cm incision is marked in the axillary crease, extending from 1 cm superior to 2 cm inferior to the lower border of the pectoralis major tendon.

fashion, and the arm is secured in a mechanical arm holder (TRIMANO Support Arm; Arthrex, Naples, FL).


#### Diagnostic Arthroscopy and LHB Tenotomy

A standard posterior portal is established, and under direct visualization, an anterosuperior portal is created. A diagnostic arthroscopy is performed. The LHB tendon, glenoid labrum, rotator cuff, chondral surfaces, and capsuloligamentous structures are evaluated. By use of a probe, the LHB tendon is pulled into the glenohumeral joint to evaluate the tendon's shape, fraying, and synovitis.

LHB tenotomy is performed with arthroscopic scissors at the superior labral insertion. A radiofrequency ablation device is then used to debride the proximal



**Fig 2.** In a right shoulder, with the pectoralis major tendon retracted superiorly (arrowhead), a right-angle clamp is used to loop around the long head of the biceps tendon (LHBT, arrow) and retrieve it from the incision. The asterisk indicates the short head of the biceps.




**Fig 3.** In a right shoulder, the periosteum in the bicipital groove (arrowhead) is stripped off with an elevator (asterisk) to prepare the cortical bed.

stump to a stable margin. Any concomitant arthroscopic procedures are carried out at this time.

#### Open Subpectoral Biceps Tenodesis With All-Suture Anchor Fixation

The complete surgical technique is shown in [Video 1](#). The arm is positioned in 90° of abduction and 90° of elbow flexion. The lower border of the pectoralis major tendon is palpated. In the axillary crease, an incision is made, extending from 1 cm superior to 2 cm inferior to the inferior border of the pectoralis major tendon ([Fig 1](#)). The pectoralis major tendon, coracobrachialis, and short head of the biceps brachii are identified. The fascia overlying the coracobrachialis and short head of the biceps is incised. With the pectoralis major tendon retracted superiorly, the bicipital groove is palpated, leading directly to the LHB tendon. A right-angle clamp is then used to loop around the LHB tendon and retrieve it from the incision ([Fig 2](#)).

The periosteum in the bicipital groove is stripped off with an elevator to prepare the cortical bed



**Fig 4.** Instruments used for all-suture anchor placement: (1) flexible 1.6-mm drill attached to arthroscopic shaver hand-piece, (2) curved drill guide, and (3) all-suture anchor with SutureTape (FiberTak).



**Fig 5.** A drill guide (arrowhead) is positioned centrally in the bicipital groove (arrow) and approximately 2 cm below the proximal edge of the pectoralis major tendon in a right shoulder. A 1.6-mm drill is used to create a unicortical bone tunnel for a single-loaded all-suture anchor (FiberTak).

(Fig 3). A drill guide is positioned centrally in the bicipital groove and approximately 2 cm below the proximal edge of the pectoralis major tendon. A 1.6-mm drill is used to create a unicortical bone tunnel for a single-loaded all-suture anchor (FiberTak) (Figs 4 and 5).

One suture strand is used to Krackow-type whipstitch the tendon with 4 passes on each side, beginning 1 cm proximal to the musculotendinous junction to a point 2 cm distal (Fig 6A). The remaining proximal portion of the tendon is sharply excised (Fig 6B). The free suture end is passed through the tendon, and applying traction on the suture reduces the tendon down to the anchor. The sutures are then tied down (Fig 7A).

The wound is irrigated and closed in a layered fashion (Fig 7B). Pearls and pitfalls of the technique, as well as advantages and disadvantages, are outlined in Tables 1 and 2.


### Postoperative Rehabilitation

For isolated subpectoral tenodesis, sling immobilization for 2 weeks with immediate active and passive range of motion is allowed. Resisted elbow flexion and forearm supination is restricted for 6 weeks after surgery. Overhead strengthening and heavy lifting are delayed for 3 months.


### Discussion

Various fixation techniques and implants are described for subpectoral biceps tenodesis.<sup>15-19</sup> Interference screw fixation has traditionally served as the gold standard with superior biomechanical strength.<sup>15</sup> However, because of concerns for a humeral stress riser with larger drill holes, other techniques have evolved to mitigate this risk while taking advantage of the benefits of the subpectoral location.

There is no clear consensus whether bone tunnel or cortical surface (onlay) healing confers better outcomes. In a rabbit model, Tan et al.<sup>20</sup> compared tendon-to-bone healing for both fixation techniques and found no significant difference between groups with respect to failure load, stiffness, and mean volume of newly formed bone. Histologic analysis showed direct tendon-to-bone healing on the outer cortical surface. In the intracortical fixation group, only 5% of the newly formed bone was located intramedullary, whereas 95% was present on the cortical surface. Siebenlist et al.<sup>21</sup> reported good clinical results and a low failure rate



**Fig 6.** Right shoulder. (A) One suture strand is used to Krackow-type whipstitch the tendon with 4 passes on each side (arrowhead), beginning 1 cm proximal to the musculotendinous junction to a point 2 cm distal. (B) The remaining proximal portion of the tendon is sharply excised (arrow).



**Fig 7.** In a right shoulder, the sutures are tied down (A) and the wound is irrigated and closed in a layered fashion (B). (LHBST, long head of biceps tendon.)

with an onlay repair technique for acute distal biceps tendon ruptures.

Several studies have reported on different supra-pectoral and subpectoral fixation techniques for biceps tenodesis, showing interference screw fixation to provide the strongest biomechanical stability.<sup>15-19,22</sup> However, complications including implant failure, bioabsorbable screw reactions, and especially humeral fractures have been reported.<sup>12-14</sup> Sears et al.<sup>14</sup> reported a case series of humeral fractures after subpectoral biceps tenodesis. They concluded that the potential stress-riser effect created by the cortical defect, location, and depth of the drill hole may be reduced by limiting the size of the cortical defect. This stress-riser effect and fracture risk may be increased especially in young overhead athletes with repetitive humeral torque.<sup>23</sup> Buchholz et al.<sup>15</sup> introduced a bone-preserving onlay technique using unicortical button fixation for subpectoral biceps tenodesis. Biomechanically, it was shown that the unicortical button withstood similar loads compared with interference screw fixation.<sup>15</sup> In a biomechanical comparison of all-suture anchor versus unicortical button fixation for subpectoral biceps tenodesis, the all-suture anchor using

a Krackow-type whipstitch provided similar biomechanical strength to unicortical button fixation using a noncontinuous locking whipstitch.<sup>24</sup> The all-suture anchor fixation technique was shown to be superior in terms of displacement during cyclic loading compared with the unicortical button fixation technique. However, the results of this study helped to show that the fixation method used on the humeral side is less implicative of overall construct strength than stitch location and technique, given that the biceps tendon tissue as well as stitch configuration seems to be the limiting factor in subpectoral onlay tenodesis techniques.

In addition to its bone-preserving approach, the all-suture anchor subpectoral biceps tenodesis is beneficial because of the aforementioned curved drilling and insertion guide, which allows for easy access to the entrance of the bicipital groove below the pectoralis major tendon, resulting in reduced surgery time. This article is a Technical Note and, as such, has a limited scope without comment on the clinical results of the described technique. With promising reported biomechanical results, further studies are needed to evaluate the clinical efficacy of this procedure.

**Table 1.** Pearls and Pitfalls of Technique

Step	Pitfalls	Pearls
Subpectoral approach	Difficulty identifying interval between pectoralis major tendon and short head of biceps	Palpate the lower border of the pectoralis major tendon.
Suture anchor placement	Incorrect suture anchor position	Position the drill guide centrally in the bicipital groove and 2 cm below the proximal edge of the pectoralis major tendon.
Tendon fixation	Inadequate tendon tensioning	Begin tendon suturing 1 cm proximal to the musculotendinous junction to a point 2 cm distal.

**Table 2.** Advantages and Disadvantages of Technique

Advantages
Predictable pain reduction with resection of proximal LHB tendon
Facilitation of adequate tendon tensioning
Reduced risk of fracture of proximal humerus
Short duration of surgery
Disadvantages
Open approach with additional scar
Potential risk of iatrogenic injury to neurovascular structures
LHB, long head of biceps.

## References

- Bain G. Biceps tendon pathology. In: Pederzini LA, Eygendaal D, Denti M, eds. *Elbow and sport*, ED 1. New York: Springer, 2016;143-152.
- Braun S, Horan MP, Elser F, Millett PJ. Lesions of the biceps pulley. *Am J Sports Med* 2011;39:790-795.
- Katthagen JC, Vap AR, Tahal DS, Horan MP, Millett PJ. Arthroscopic repair of isolated partial- and full-thickness upper third subscapularis tendon tears: Minimum 2-year outcomes after single-anchor repair and biceps tenodesis. *Arthroscopy* 2017;33:1286-1293.
- Pogorzelski J, Horan MP, Hussain ZB, Vap A, Fritz EM, Millett PJ. Subpectoral biceps tenodesis for treatment of isolated type II SLAP lesions in a young and active population. *Arthroscopy* 2018;34:371-376.
- Provencher MT, LeClere LE, Romeo AA. Subpectoral biceps tenodesis. *Sports Med Arthrosc Rev* 2008;16:170-176.
- Longo UG, Loppini M, Marineo G, Khan WS, Maffulli N, Denaro V. Tendinopathy of the tendon of the long head of the biceps. *Sports Med Arthrosc Rev* 2011;19:321-332.
- Hsu AR, Ghodadra NS, Provencher MT, Lewis PB, Bach BR. Biceps tenotomy versus tenodesis: A review of clinical outcomes and biomechanical results. *J Shoulder Elbow Surg* 2011;20:326-332.
- Slenker NR, Lawson K, Ciccotti MG, Dodson CC, Cohen SB. Biceps tenotomy versus tenodesis: Clinical outcomes. *Arthroscopy* 2012;28:576-582.
- Vap AR, Katthagen JC, Tahal DS, et al. Isolated biceps reflection pulley tears treated with subpectoral biceps tenodesis: Minimum 2-year outcomes. *Arthroscopy* 2017;33:1788-1794.
- Millett PJ, Sanders B, Gobeze R, Braun S, Warner JJ. Interference screw vs. suture anchor fixation for open subpectoral biceps tenodesis: Does it matter? *BMC Musculoskelet Disord* 2008;9:121.
- Park JS, Kim SH, Jung HJ, Lee YH, Oh JH. A prospective randomized study comparing the interference screw and suture anchor techniques for biceps tenodesis. *Am J Sports Med* 2017;45:440-448.
- Euler SA, Smith SD, Williams BT, Dornan GJ, Millett PJ, Wijdicks CA. Biomechanical analysis of subpectoral biceps tenodesis: Effect of screw malpositioning on proximal humeral strength. *Am J Sports Med* 2015;43:69-74.
- Nho SJ, Reiff SN, Verma NN, Slabaugh MA, Mazzocca AD, Romeo AA. Complications associated with subpectoral biceps tenodesis: Low rates of incidence following surgery. *J Shoulder Elbow Surg* 2010;19:764-768.
- Sears BW, Spencer EE, Getz CL. Humeral fracture following subpectoral biceps tenodesis in 2 active, healthy patients. *J Shoulder Elbow Surg* 2011;20:e7-e11.
- Buchholz A, Martetschlager F, Siebenlist S, et al. Biomechanical comparison of intramedullary cortical button fixation and interference screw technique for subpectoral biceps tenodesis. *Arthroscopy* 2013;29:845-853.
- Altintas B, Pitta R, Fritz EM, Higgins B, Millett PJ. Technique for type IV SLAP lesion repair. *Arthrosc Tech* 2018;7:e337-e342.
- Chiang FL, Hong CK, Chang CH, Lin CL, Jou IM, Su WR. Biomechanical comparison of all-suture anchor fixation and interference screw technique for subpectoral biceps tenodesis. *Arthroscopy* 2016;32:1247-1252.
- Kennedy NI, Sanchez G, Mannava S, Ferrari MB, Frangiamore SJ, Provencher MT. Arthroscopic rotator cuff repair with mini-open subpectoral biceps tenodesis. *Arthrosc Tech* 2017;6:e1667-e1674.
- Su WR, Ling FY, Hong CK, Chang CH, Lin CL, Jou IM. Subpectoral biceps tenodesis: A new technique using an all-suture anchor fixation. *Knee Surg Sports Traumatol Arthrosc* 2015;23:596-599.
- Tan H, Wang D, Lebaschi AH, et al. Comparison of bone tunnel and cortical surface tendon-to-bone healing in a rabbit model of biceps tenodesis. *J Bone Joint Surg Am* 2018;100:479-486.
- Siebenlist S, Schmitt A, Imhoff AB, et al. Intramedullary cortical button repair for distal biceps tendon rupture: A single-center experience. *J Hand Surg Am* 2019;44:418e1-418.e7.
- Kennedy NI, Godin JA, Ferrari MB, et al. Subpectoral biceps tenodesis: Interference screw and cortical button fixation. *Arthrosc Tech* 2017;6:e1415-e1420.
- Sabick MB, Torry MR, Kim YK, Hawkins RJ. Humeral torque in professional baseball pitchers. *Am J Sports Med* 2004;32:892-898.
- Lacheta L, Rosenberg SI, Brady AW, Dornan GJ, Millett PJ. Biomechanical comparison of subpectoral biceps tenodesis onlay techniques. *Orthop J Sports Med* 2019;7:2325967119876276.